

# Statistics for Astronomers

## Homework #8 (Due before 12:00 PM on Monday, 2021.01.25)

Prof. Sundar Srinivasan

January 17, 2021

**Notes:** (1) You are welcome to use `Python` functions to evaluate probabilities for various distributions, and `Mathematica/Wolfram Alpha` to compute integrals if necessary. **Just mention your source in each case!** (2) Email me your `Python` scripts and any/all resulting output plots/images.

1. The flux density  $F_{\nu, \text{BB}}(T_{\text{eff}}, R_{\text{eff}}, D)$  in mJy at wavelength  $\lambda = 7.872 \mu\text{m}$  received from a blackbody is given by

$$F_{\nu, \text{BB}}(T_{\text{eff}}, R_{\text{eff}}, D) = \frac{0.26184 \left( \frac{R_{\text{eff}}}{D} \right)^2}{\exp \left( \frac{1848.9}{T_{\text{eff}}} \right) - 1},$$

with  $R_{\text{eff}}$  in AU,  $D$  in kpc, and  $T_{\text{eff}}$  in K. Assume that the values for  $(R_{\text{eff}}, D, \text{ and } T_{\text{eff}})$  are normally distributed about means (1.330 AU, 0.892 kpc, 3393 K) with standard deviations equal to (0.300 AU, 0.092 kpc, 275 K).

- (a) **(3 points)** Find the mean, median, and mode for the distribution of fluxes generated by propagating the uncertainties in the radius, distance, and temperature.
  - (b) **(3 points)** What are the upper and lower limits for the 68% equal-tailed interval?
2. Read through the procedure for rejection sampling described here. In this problem, you will use rejection sampling to draw random variables from the blackbody distribution (this will be our **target distribution**)

$$p(x) = \frac{15}{\pi^4} \frac{x^3}{e^x - 1}.$$

The **proposal distribution** for this problem will be a (truncated) normal distribution.

- (a) **(2 points)** Choose appropriate values of the mean and standard deviation for the proposal distribution.
- (b) **(2 points)** Choose an appropriate scaling factor to multiply into the proposal distribution so that it is greater than or equal to the target distribution for all values of  $x$ .

- (c) **(2 points)** Over the relevant range of  $x$  values, what fraction of the area under the proposal distribution is also covered by the target distribution? Based on this, what is the expected value for the fraction of points that will be rejected?
- (d) **(6 points)** Starting with  $N = 10\,000$  draws from the proposal distribution, write a script that uses rejection sampling to simulate draws from the target distribution.
- (e) **(4 points)** As part of the same script, generate a kernel density estimate (you can use `statsmodels.nonparametric.kde.KDEUnivariate`) from the simulated draws. Compare this to the target distribution function on a plot to verify that it has the proper shape.