

The nature of faint radio galaxies at high redshifts

A. Saxena^{1b, 1,2★}, H. J. A. Röttgering¹, K. J. Duncan^{1b, 1}, G. J. Hill^{3,4}, P. N. Best⁵,
B. L. Indahl³, M. Marinello⁶, R. A. Overzier^{6,7}, L. Pentericci², I. Prandoni^{1b, 8}, H. Dannerbauer^{9,10} and
R. Barrena^{9,10}

¹Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands

²INAF-Osservatorio Astronomico di Roma, Via Frascati 33, I-00040 Monteporzio (RM), Italy

³Department of Astronomy, University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712, USA

⁴McDonald Observatory, University of Texas at Austin, 2515 Speedway, Stop C1402, Austin, TX 78712, USA

⁵Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

⁶Observatório Nacional, Rua General José Cristino, 77, São Cristóvão, Rio de Janeiro, RJ, CEP 20921-400, Brazil

⁷Institute of Astronomy, Geophysics and Atmospheric Sciences, Department of Astronomy, University of São Paulo, São Paulo, SP 05508-090, Brazil

⁸INAF-Istituto di Radioastronomia, Via P. Gobetti 101, I-40129 Bologna, Italy

⁹Instituto de Astrofísica de Canarias (IAC), E-38205 La Laguna, Tenerife, Spain

¹⁰Universidad de La Laguna, Dpto. Astrofísica, E-38206 La Laguna, Tenerife, Spain

Accepted 2019 September 4. Received 2019 September 4; in original form 2019 June 3

ABSTRACT

We present spectra and near-infrared images of a sample of faint radio sources initially selected as promising high-redshift radio galaxy (HzRG) candidates. We have determined redshifts for a total of 13 radio galaxies with redshifts ranging from $0.52 \leq z \leq 5.72$. Our sample probes radio luminosities that are almost an order of magnitude fainter than previous large samples at the highest redshifts. We use near-infrared photometry for a subsample of these galaxies to calculate stellar masses using simple stellar population models, and find stellar masses to be in the range $10^{10.8} - 10^{11.7} M_{\odot}$. We then compare our faint radio galaxies with brighter radio galaxies at $z \geq 2$ from the literature. We find that fainter radio galaxies have lower $\text{Ly}\alpha$ luminosities and narrower line widths compared to the bright ones, implying photoionization by a weaker active galactic nucleus (AGN). We also rule out the presence of strong shocks in faint HzRGs. The stellar masses determined for faint HzRGs are lower than those observed for brighter ones. We find that faint HzRG population in the redshift range 2–4 forms a bridge between star-forming and narrow-line AGNs, whereas the ones at $z > 4$ are likely to be dominated by star formation, and may be building up their stellar mass through cold accretion of gas. Finally, we show that the overall redshift evolution of radio sizes at $z > 2$ is fully compatible with increased inverse Compton scattering losses at high redshifts.

Key words: galaxies: active – galaxies: high-redshift.

1 INTRODUCTION

High-redshift radio galaxies (HzRGs) have been found to be hosted by some of the most massive galaxies in the Universe and are thought to be progenitors of the massive ellipticals we observe today. HzRGs contain large amounts of dust and gas (Best et al. 1998b; Archibald et al. 2001; Carilli et al. 2002; Reuland et al. 2004; De Breuck et al. 2010), are dominated by an old stellar population (Best, Longair & Röttgering 1998a; Rocca-Volmerange et al. 2004; Seymour et al. 2007), and seen to have high star formation rates (Willott et al. 2003; McLure et al. 2004; Miley et al. 2006; Villar-Martín et al. 2007a; Seymour et al. 2008). Owing to the large stellar masses of their host galaxies, radio-loud active galactic nuclei (AGNs) across all redshifts are often found to be located in the centre of clusters and proto-clusters (Hill & Lilly 1991; Pentericci et al. 2000; Venemans et al. 2002; Röttgering et al. 2003; Miley et al. 2004; Hatch et al. 2011;

Wylezalek et al. 2013; Orsi et al. 2016). Miley & De Breuck (2008) summarize the properties of radio galaxies and their environments in their review.

HzRGs are seen to be some of the most massive galaxies across all redshifts (see Rocca-Volmerange et al. 2004, for example). As a result, near-infrared (NIR) magnitudes of radio galaxies have been observed to follow a correlation with redshift, which is known as the K - z relation (Jarvis et al. 2001b; Willott et al. 2003; Rocca-Volmerange et al. 2004). This is mainly because of the consistently high stellar masses observed and because HzRGs are thought to be dominated by an old stellar population, and the K band traces the part of spectral energy distribution (SED) that contains light emitted from these old stars. At higher redshifts ($z > 4$), the scatter in the K - z relation is seen to increase as bluer parts of the rest-frame SED containing light from younger stars (which is more sensitive to the star formation history) are observed in the K band.

Selecting HzRGs based on their radio emission alone has the advantage that no prior colour selection (one that probes the Lyman break, for example) is applied (De Breuck et al. 2000a). In principle,

* E-mail: saxena@strw.leidenuniv.nl

promising high-redshift candidates selected from a radio survey with follow-up optical/IR spectroscopy could discover massive galaxies all the way into the epoch of reionization (EoR) through observations of their redshifted synchrotron spectrum. Since radio observations are not affected by dust, radio selected galaxies can also provide an unbiased sample to study the dust properties of galaxies at high redshifts.

A successful method of identification of HzRGs from all-sky radio surveys is to first create samples of radio sources that display an ultra-steep spectral index (α , where $S \propto \nu^\alpha$) and then follow these up at optical and infrared wavelengths (Tielens, Miley & Willis 1979; Röttgering et al. 1994; Chambers et al. 1996; Blundell, Rawlings & Willott 1999; De Breuck et al. 2000a; Ishwara-Chandra et al. 2010; Afonso et al. 2011; Singh et al. 2014). This method is effective owing to the apparent correlation that exists between the redshift of a radio source and the steepness of its low-frequency radio spectrum (see Ker et al. 2012, for example). The physical explanation for this correlation is still under debate, but it is commonly accepted that curvature of the radio SED combined with a K -correction is responsible for the apparent correlation. For a more thorough discussion on the redshift–spectral index correlation, we point the readers to Miley & De Breuck (2008) where some other possible explanations are also discussed. From an observational point of view, however, radio sources with an ultra-steep spectrum (USS) generally make promising HzRG candidates. Most of the currently known radio galaxies at the highest redshifts have been selected from relatively shallow all-sky surveys. Therefore, the well-studied HzRG samples (van Breugel et al. 1999; Willott et al. 1999; De Breuck et al. 2000a; Jarvis et al. 2001a; Rawlings, Eales & Lacy 2001; Willott et al. 2003; Vardoulaki et al. 2010; Rigby et al. 2011) mostly consist of luminous radio sources, and very little is currently known about the fainter and presumably more abundant radio galaxies at high redshifts. With the advent of the latest generation of radio telescopes such as the upgraded Giant Metrewave Radio Telescope (GMRT; Swarup et al. 1991), the Low-Frequency Array (LOFAR; van Haarlem et al. 2013), the Murchinson Widefield Array (MWA; Tingay et al. 2013), and the Square Kilometer Array (SKA; Dewdney et al. 2009) in the future, it is now possible to survey large areas of the sky to unprecedented depths at low radio frequencies and push the discovery of radio galaxies to lower luminosities and/or to higher redshifts, well into the EoR ($z > 6$).

The spectra of radio galaxies across all redshifts are rich with emission lines and there have been many studies exploring the primary source of photoionization in radio galaxies and AGNs using UV and optical emission lines and their ratios. Robinson et al. (1987) and Baum, Heckman & van Breugel (1992) found that photoionization by the central AGN alone was found to sufficiently explain the UV and optical line ratios observed in low-redshift radio galaxies ($z < 0.2$). However, for radio galaxies at higher redshifts, photoionization by an AGN was not enough to explain the observed blue colours and bright Ly α emission observed in these objects (Villar-Martin, Tadhunter & Clark 1997). Further, there is evidence of strong interaction between the radio jet and the interstellar medium (ISM) of the host galaxy, which leads to the generation of powerful shocks that disturb the morphology, kinematics, and physical conditions of the line-emitting gas (McCarthy, Baum & Spinrad 1996; van Ojik et al. 1996; Villar-Martin et al. 1997). This interaction is also supported by observations that show close alignment between the radio jet and the extended emission-line regions (EELRs) of HzRGs (Miley et al. 1992; Jannuzi et al. 1995). More recently, observations of the alignment of molecular gas with the radio jet in distant radio galaxies have lent further support to the

jet–gas interaction scenario in radio galaxies (see Emonts et al. 2014, for example).

In this paper, we first present results from a campaign to spectroscopically follow-up the faint HzRG candidates identified by Saxena et al. (2018a) and observe a subset of these at NIR wavelengths. The layout of this paper is as follows. In Section 2, we present our new sample of spectroscopically confirmed radio galaxies. We first briefly describe the sample selection, observations, and data reduction methods. We then present the spectroscopic and galaxy properties derived for the new radio galaxy sample, which probes fainter flux densities than previous all-sky radio surveys across all redshifts. In Section 3, we compile a list of known radio galaxies at $z > 2$ from the literature, and use this literature sample to compare the properties of the faint high-redshift radio galaxies presented in this paper. We explore various correlations between radio properties such as luminosity, size, and spectral index with emission line properties such as line luminosity and widths in both the faint and literature HzRG samples. In Section 4, we present a discussion about the possible nature of faint HzRGs and offer an explanation for the overall trends observed in the radio galaxy samples. Finally, in Section 5 we summarize the key findings of this study.

Throughout this paper, we assume a flat lambda cold dark matter cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_m = 0.3$.

2 THE FAINT HZRG SAMPLE

2.1 Sample selection

The HzRG candidates targeted for spectroscopy are mainly selected on the basis of (i) steepness of their spectral index ($\alpha_{1400}^{150} < -1.3$) calculated by cross-matching with the Very Large Array (VLA) FIRST survey at 1.4 GHz (Becker, White & Helfand 1995) and (ii) compact morphologies (largest angular size < 10 arcsec), which are expected from radio galaxies at the highest redshifts (Saxena, Röttgering & Rigby 2017). Lower flux density limits were applied both at 150 MHz and 1.4 GHz to ensure a high enough signal-to-noise ratio and upper flux density limits were applied at 150 MHz to ensure that none of the sources in our final sample would have been detected in previous searches for USS radio sources (for example, De Breuck et al. 2000a). Further, only those sources with no clear optical or infrared counterparts in the available all-sky surveys such as SDSS (Alam et al. 2015), PanSTARRS1 (Chambers et al. 2016), ALLWISE (Wright et al. 2010), and UKIDSS LAS (Lawrence et al. 2007) were retained in the final sample, increasing the likelihood of a high-redshift nature of the radio host. The distribution of the selected sources in the radio flux density – spectral index parameter space (taken from Saxena et al. 2018a) – is shown in Fig. 1.

The final sample consisted of 32 radio sources that were followed up with the Karl G. Jansky VLA in configuration A in bands L and P (Project code: 16B-309, PI: Röttgering), with an average beam size of 1.3 arcsec at 1.4 GHz (L band). These results are presented in Saxena et al. (2018a). The high-resolution VLA images enabled subarcsecond isolation of the expected optical counterpart of the radio source. Blind spectroscopy at the position of the expected host of a radio source has been found to be successful at detecting emission lines and obtaining redshifts (Röttgering et al. 1994; De Breuck et al. 2001).

2.2 Observations and data reduction

Spectra presented in this paper were obtained over a period of 2016 December–2019 March using three different telescopes, namely the




Figure 1. The unique parameter space in flux density and spectral index probed by the Saxena et al. (2018a) sample of USS radio sources selected from TGSS ADR1. Shown for comparison are the radio sources from De Breuck et al. (2000a). Going to fainter flux densities should open up the potential for discovering radio galaxies at higher redshifts than seen before, and lead to the discovery of fainter radio galaxies at all redshifts, providing a more complete picture of the radio galaxy population.

10 m Hobby–Eberly Telescope (HET) at McDonald Observatory, Texas, the 8.2 m Gemini North at Hawaii, and the 4.2 m William Herschel Telescope (WHT) at La Palma, Canary Islands. Due to the large distribution in right ascension of the sample, a homogeneous follow-up strategy was not practical and spectra had to be obtained over a long period of time, through multiple observing runs on numerous facilities around the world.

We used the long-slit set-up with Gemini and WHT, and the integral field unit (IFU) set-up with the HET. The typical observing times were 1 h on source per object. The radio morphologies for most of the targets were unresolved (at a beam size of ~ 1.3 arcsec, and for these targets we aligned the slit at a parallactic angle of 0 deg. For resolved radio objects, the slit was aligned along the extended radio morphology, as radio galaxy hosts are typically extended over a few arcseconds and often oriented along the radio axis (Röttgering et al. 1997). IFU observations were simply centred on the expected location of the radio host galaxy. Below we describe the observations taken using each telescope in detail. An overview of the set-ups used in various observing runs is given in Table 1.

2.2.1 4.2 m WHT

The observations at the WHT were carried out using the Intermediate dispersion Spectrograph and Imaging System (ISIS¹) instrument with a 3 arcmin slit length and a slit width of 1.5 arcsec (Program IDs: W17AN014 and W19AN015; PI: Saxena). We used the R300B grating in the Blue arm and the R158R grating in the Red arm of the spectrograph, providing a spectral resolution of $R \sim 975$ and 910, respectively, and a total spectral range of $(\sim 3200) - (\sim 9200)$ Å over both arms. The first set of observations is carried out from 2017 July 24 to 27. The first two nights of the observing run were strongly affected by Saharan dust covering the island, leading to extinction of up to 2 mag in worst conditions. The conditions improved significantly for the final two nights. The average seeing over all four nights was around 0.6 arcsec. The second set of observations

is carried out from 2019 March 3 to 5. The average seeing for the first two nights was ~ 1.0 arcsec, and the third night was lost due to bad weather. During both runs, the same strategy for calibration was adopted, as described below.

Standard afternoon calibrations were performed and both dome and sky flats were taken before and after each observing night. A flux standard was observed at the beginning and the end of each night. Each target was observed for a total of 3600 (1800×2) s in each arm simultaneously, except when a bright line or continuum was spotted after the first 1800 s. In such cases, the second exposure was terminated earlier. We used CuAr + CuNe lamps for wavelength calibration, which were observed at the position of each target before the sky exposure. Since the optical magnitudes of the targets are very faint ($R > 23.5$ AB), we first acquired a bright star close to the target and then added offsets to arrive at the target coordinates. This ensures accurate positioning of the slit for very faint targets. Standard data reduction procedures that include bias subtraction, flat-fielding, sky subtraction, wavelength calibration, and flux calibration were performed using PYTHON scripts written by our team. These scripts were based on the WHT data reduction package written by Steve Crawford.²

2.2.2 8.2 m Gemini North

The observations at Gemini North were carried out using the Gemini Multi-Object Spectrograph (GMOS; Hook et al. 2004) in long-slit mode with a slit width of 1.5 arcsec and a slit length of 5 arcmin (Program ID: GN-2017A-Q-8; PI: Overzier). A total observing time of 7 h was used to observe six targets. The observations were carried out over a period of 2 months, from 2017 April 7 to 2017 June 24. We used the R400_G5305 grating with a central wavelength of 700 nm, giving a resolution of $R \sim 1500$, providing coverage over the wavelength range 3600–9400 Å. We took three exposures of 800 s each, giving 2400 s of exposure time per object.

We used the Gemini IRAF package³ for reducing the data, which performed all the basic reductions including bias subtraction, dark subtraction, flat-fielding, and cosmic ray removal. Wavelength and flux calibration was then performed on the spectra using the pipeline tools.

2.2.3 10 m HET

The observations at the HET were carried out using the Low-Resolution Spectrograph 2 (LRS2; Chonis et al. 2016), which is a 6×12 sq. arcsec lenslet-coupled fibre IFU. The wavelength coverage of the LRS2 is split into two spectrographs, the LRS2-B, which covers 3650–4670 Å and 4540–7000 Å at resolving powers of $R \approx 2500$ and 1400, respectively, and the LRS2-R, which covers 6430–8450 Å and 8230–10 560 Å at resolving powers of $R \approx 2500$ each. The observations were taken in spectroscopic sky conditions over four trimesters, beginning 2016 December and ending 2018 March (PI: Hill).

We used the LRS2 quicklook reduction pipeline⁴ written in PYTHON (Indahl et al. submitted) to perform the basic reduction of the IFU data. The reduction steps include bad pixel masking, bias subtraction, flat-fielding, cosmic ray removal, and wavelength calibration using arc lamps (taken with a combination of Cd, Hg,

²Can be found at: https://github.com/crawfordsm/wht_reduction_scripts.

³<https://www.gemini.edu/sciops/data-and-results/processing-software>

⁴https://github.com/BrianaLane/LRS2_reduction

¹<http://www.ing.iac.es/astronomy/instruments/isis/index.html>

Table 1. Observational set-ups for spectroscopy used in this study.

Telescope	Instr.	Grism/Grating	λ coverage (Å)	Date	Exp. time (sec)	Limiting flux (1σ) (erg s ⁻¹ cm ⁻² Å ⁻¹)
WHT	ISIS	R300B/R158R	3200–9200	2017 Jul. 24–27	1800–3600	1.1×10^{-18}
–	–	–	–	2019 Mar. 3–5	3600–7200	2.7×10^{-18}
Gemini North	GMOS	R400-G5305	5600–9400	2017 Apr. 7–Jun. 24	2400	1.0×10^{-18}
HET	LRS2 IFU	LRS2-B/R	3650–10560	2016 Dec.–2018 Mar.	3600	$2.0\text{--}7.0 \times 10^{-18}$

FeAr, or Kr, covering the full wavelength range) for individual frames from both blue and red arms. The reduced data products include both 2D row-stacked spectra (RSSs) and 3D data cubes. The individual RSS and cubes were then median combined to form the final science data products.

The presence of line emission in the data was first identified by visually examining the 2D spectra from each arm of the spectrograph. The 2D spectrum generated for each arm by the data reduction pipeline is essentially an RSS of all the IFU fibres. Presence of an emission line in an arm can be distinguished from any other spurious sources of flux by searching for signs of emission across multiple fibres. This is shown in the left-hand side panels of Fig. 2 for two sources. USS18 (top-left) has a strong emission line, and USS46 (bottom-left) has a weaker emission line. The distribution of emission-line flux over multiple fibres helps rule out cosmic rays or hot pixels and confirm that the emission is real. The spatial element (lenslet) size of the LRS2 IFUs is 0.6 arcsec, so typical objects fall on several individual lenslets/fibres depending on the image quality of the observation. The spectra resulting from those fibres are separate on the CCD. Additionally, light from an unresolved emission line on each fibre falls on 14 pixels, more if the line is resolved. The combination of these factors results in a distinct pattern of separated, resolved spectra which is impossible to confuse with a CR hit (as shown in Fig. 2). Single-line detections with LRS2 are more secure than for a long-slit spectrograph in this respect.

Where signs of an emission line were spotted in the 2D spectrum, we looked at the spatial distribution of the line emission in the 3D data cube generated by the pipeline for that arm of the spectrograph. This is shown in the right-hand side panels of Fig. 2 for both a strong and a weak emission line. A circular aperture that fully encompassed pixels with a signal-to-noise of 3 or above was then used to extract a 1D spectrum of the source. The LRS2-B and -R response functions were used to calibrate the flux of the extracted 1D spectrum and emission lines were then identified using the methods that are described in the following section.

For sources with an emission line in only one arm of the spectrograph, we could not extract spectra from the other arms where no signs of emission lines were found. This is primarily because the quicklook data reduction pipeline for HET-LRS2 is unable to assign a World Coordinate System (WCS) to the individual fibres of each arm of the spectrograph, and therefore, in the absence of an emission line it is not possible to extract a spectrum by placing an aperture on a particular spatial position on the sky. A future updated pipeline will enable extraction of 1D spectra for all sources, regardless of whether or not an emission-line feature has been spotted. However, since the field of view of LRS2 is relatively small (6 arcsec $\times$ 12 arcsec), any detection of an emission-line source is likely to be the counterpart of the radio source that has been targeted.

Additionally, the pointing accuracy of HET is demonstrated by the detection of emission lines from sources that also have spectra from either WHT or Gemini, and by pointing tests on stars. Although we attempted to observe different targets in each of our spectroscopy runs

on the three telescopes listed above to increase the number of radio sources with redshifts, there were a few targets that were observed using multiple telescopes owing to target visibility and/or scheduling restrictions. Consistent emission-line features were observed across observations using different facilities, which further helped confirm that the emission lines are real. Although specifically for the HET data cubes (with a field of view of 6 arcsec $\times$ 12 arcsec, there may be an element of uncertainty associated with whether the line-emitting component is actually a part of the radio galaxy. The initial radio source selection, however, takes care to eliminate any possible low-redshift interlopers.

We list the number of targets observed and the number of detections, either emission lines or continuum, recorded during all our runs in Table 2. In the next section, we summarize the flux limits achieved and detection rates from each of our observing runs.

2.2.4 NIR imaging

Additional NIR imaging in the *J* and *K* bands was obtained using LUCI (formerly known as LUCIFER; Seifert et al. 2003) on the Large Binocular Telescope (LBT) for nine HzRG candidates in our sample. Of these, seven had spectroscopically confirmed redshifts. Below we highlight the observations and data reduction strategy.

The observations were carried out in two separate runs, with the first on 2018 February 1 and the second on 2018 May 11 (Program ID: 2017 2018 43; PI: Prandoni). The average seeing throughout the observations was 0.6–1.0 arcsec. The average on-source exposure times were 720 (12 $\times$ 60 s) seconds in *J* (central wavelength of 1.247 microns) and 1200 (20 $\times$ 60 s) in *K* (central wavelength of 2.194 microns). We used a standard jitter pattern with a maximum jitter distance of 15 arcsec to enable effective sky subtraction in both the bands.

The LUCI data reduction pipeline developed at INAF-OAR was used to perform the reductions and calibrations of raw images. These tasks include dark subtraction, bad pixel masking, cosmic ray removal, flat-fielding, and sky subtraction. Astrometric corrections for individual frames were calculated and applied and the corrected frames were then resampled and combined using a weighted co-addition method to obtain a deep image. The photometry of the co-added image was then calibrated using 2MASS and where available, UKIDSS Large-Area Survey to an accuracy of within 0.1 mag. The median and standard deviation of the background in all co-added images were then calculated by placing random apertures with a diameter of 1.5 arcsec.

One source was observed using EMIR, a NIR imager/spectrograph (Garzón et al. 2016) on the Gran Telescopio Canarias (GTC) in queue mode (Proposal ID: 124-GTC85/17B, PI: Génova Santos). The observation was taken in the *J* and *K* bands, with 220 and 760 s of on-source exposure time, respectively. The average seeing for this observation was 1.0 arcsec. Standard procedures for data reduction and calibrations of the raw images were performed, as mentioned previously.




Figure 2. A visual representation of emission line identification and 1D spectrum extraction from the HET IFU data. The left-hand side panels show the 2D RSS from the individual fibres of HET-LRS2 for a strong (top) and a weak (bottom) line-emitting source, and the right-hand side panels show the spatial distribution of the emission line in a slice from the 3D cube corresponding to the wavelength where the emission-line flux is at its peak. The 2D spectra were used to search for emission lines, and the 3D cubes were then used to extract the 1D spectrum using a circular aperture.

Table 2. The number of objects observed and the number of detections during spectroscopy runs on different telescopes. There are a few objects that have duplicate observations on multiple telescopes. We also count continuum detections in these numbers, but their spectra have not been presented in this study.

Telescope (code)	Observed	Detected	Percentage detected
WHT (W)	20	8	40 per cent
Gemini North (G)	6	5	83 per cent
HET (H)	15	8	54 per cent

2.3 Results

2.3.1 Flux limits and detection rates

The median 1σ flux limit achieved in the first observing run on WHT was $1.1 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. We targeted 12 sources in total and were able to detect 7 (either emission lines or continuum). During the second observing run on the WHT, the conditions were inferior compared to the first run and the median flux limit achieved was $2.7 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. We observed nine sources in total and detected an emission line in only one source. No continuum was detected in any other source either.

The flux limits reached during observations with the HET are more variable due to queue scheduling and the variable illumination during tracking. We find for our observations, which were taken across

different seasons, 1σ noise levels ranging from $2.0\text{--}7.0 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. On average, the blue channel is more sensitive compared to the red channel due to the lower sky background. The average flux limits reached during six different observations using Gemini North are $1.0 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. The flux limits are also shown in Table 1.

2.3.2 Redshift determination and emission line measurements

After applying all the calibrations, the identification of emission lines in the extracted 1D spectra was attempted. The spectra of radio galaxies are dominated by one or several of the following emission lines: Ly α λ 1216, [O II] λ 3727, [O III] λ 5007, H α λ 6563 + [N II] λ 6583, and the initial redshift determination was based on the identification of one or more of these lines (McCarthy 1993; Röttgering et al. 1997; De Breuck et al. 2001). Relatively fainter lines such as C IV λ 1549, He II λ 1640, C III] λ 1909, C II] λ 2326, and Mg II λ 2800 were then used in case of ambiguity arising from the identification of the stronger emission lines. For objects with a single emission line in the spectrum, we used the shape and the equivalent width of the emission line, in addition to NIR magnitudes (where available) to identify the emission line for redshift determination. For example, an asymmetric line profile (combined with a faint NIR magnitude) would be a strong indicator of Ly α at high redshifts.

Once the redshift was determined, we derived the integrated flux and full width at half-maximum (FWHM), by fitting a single Gaussian to the emission line. In a few cases, we detect double-peaked emission lines (mostly Ly α λ 1216 and [O II] λ 3727), but for the purposes of this study we smoothed the spectrum in order to fit a single Gaussian to such lines. The above line identification methods yielded reliable redshifts for a total of 12 objects, which are presented in Table 3. Also shown are the flux densities and calculated radio luminosities at 150 MHz from Saxena et al. (2018a). The 1D spectra are shown in Fig. A1 in Appendix A.

2.3.3 Notes on individual object spectra

(i) USS7, $z = 0.53$

Two narrow emission lines are seen in the spectrum, which we identify as [O III] λ 5007 and [O II] λ 3727 at a redshift of $z = 0.53$. Narrow [Ne III] λ 3967 is potentially also present, although it lies in a region of strong sky lines. This source is slightly resolved in the radio image and the [O III] and [O II] line strengths suggest that this source is a faint high-excitation radio galaxy.⁵

(ii) USS43, $z = 1.43$

A redshift of $z = 1.43$ was confirmed using the [O II] λ 3727 and Mg II λ 2800 emission lines. Also detected in the spectrum is [Ne V] λ 3426, which has been shown to be indicative of the presence of EELRs due to the AGN in the host galaxy, which also contribute towards powering the [O II] emission in addition to star formation activity (Maddox 2018). When applying a smoothing kernel to the spectrum, we also see faint C II] λ 2324, C III] λ 1909, and C IV λ 1549 emission. C III] is blue-shifted by 30 Å, which is a better indicator of the systemic redshift of the galaxy. However, since the signal-to-noise of this line is low, we quote the redshift measured using other lines instead. Our NIR imaging led to the detection of the host galaxy with $K = 21.6 \pm 0.2$ AB.

(iii) USS483, $z = 1.49$

This object has an interesting spectrum with only one bright emission line visible at 9291 Å. There is faint continuum visible blueward of the emission line and the line appears to be symmetric, therefore ruling out the line identification as Ly α at $z \sim 6.6$ where we would expect to see an asymmetric line profile and no continuum blueward of the line.

We tentatively identify this line as [O II] λ 3727 because of the slight hint of a double-peaked morphology, giving a redshift of $z = 1.49$. There is very faint emission (SNR < 2) visible in the spectrum that is coincident with Ne V at this redshift. H α is ruled out due to the absence of the expected [O III] line in this case, and [O III] is also ruled out since we would expect to see [O II] λ 3727 in the blue part of the spectrum. The redshift of this source remains uncertain and a deeper spectrum is required to uncover other emission lines that will help confirm the spectroscopic redshift.

The long-slit was aligned along the radio axis (Fig. 3, top) and the line emission in the 2D spectrum (Fig. 3, bottom) is extended along the radio axis, which implies interaction between the jet and the gas surrounding the host galaxy. The line emission can be seen extending up to 15 arcsec, pointing towards the possibility of large-scale emission-line haloes around the central radio galaxy.

⁵High-excitation radio galaxies (HERGs) are a class of radio galaxies that show strong high-excitation emission lines. A characteristic feature of HERGs is their high [O III] equivalent widths (> 3 Å) and high [O III]/[O II] ratios (see Best & Heckman 2012, for example).

(iv) USS182, $z = 2.02$

A strong emission line is detected in the spectrum. No other lines or continuum is detected. We identify this line as Ly α at $z = 2.02$ as the line is asymmetric, and lies at a wavelength that is too short for it to be [O II] or any of the other bright lines typically observed in the spectrum of radio galaxies. The emission line has an integrated flux of $32.6 \pm 5.0 \times 10^{-17}$ erg s⁻¹ cm⁻² and a broad FWHM of 1420 ± 60 km s⁻¹. Our NIR imaging led to the detection of the host galaxy with $K = 21.8 \pm 0.2$ AB.

(v) USS18, $z = 2.06$

A strong emission line, which we identify as Ly α at $z = 2.06$, is detected in the spectrum. No other lines or continuum is detected. This line is identified as Ly α because of the same reasons as for USS182 – an asymmetric line profile and too short a wavelength for [O II]. The emission line has an integrated flux of $77.9 \pm 4.3 \times 10^{-17}$ erg s⁻¹ cm⁻² and an FWHM of 950 ± 60 km s⁻¹.

(vi) USS320, $z = 2.20$

A double-peaked emission line is observed in the spectrum, which we identify as Ly α at $z = 2.20$. The line has a characteristic double-peaked shape with H I absorption between the two peaks, often associated with Ly α from radio galaxies (see van Ojik et al. 1997, for example). There are also hints of very faint C IV and possible Mg II emission (although in sky lines) in the redder parts of the spectrum at the same redshift. We smooth the double-peaked Ly α emission line and fit with a single Gaussian, deriving a line flux of $8.7 \pm 3.8 \times 10^{-17}$ erg s⁻¹ cm⁻² and an FWHM of 1300 ± 100 km s⁻¹. The host galaxy is detected with a magnitude of $K = 21.7 \pm 0.3$ AB in our NIR imaging.

(vii) USS268, $z = 2.35$

A double-peaked emission line is seen in the spectrum, which we identify as Ly α at $z = 2.35$. The line is identified as Ly α for the same reasons as the source above – double-peaked profile with H I absorption between the peaks. Mg II is also detected in the redder part of the spectrum at the same redshift, although this is in a region affected by strong sky lines. We smooth the double-peaked Ly α line and fit with a single Gaussian to calculate a line flux of $22.8 \pm 2.8 \times 10^{-17}$ erg s⁻¹ cm⁻² and an FWHM of 750 ± 50 km s⁻¹. Our NIR imaging led to the detection of the host galaxy with $J = 23.6 \pm 0.1$ AB and $K = 21.9 \pm 0.3$ AB.

(viii) USS98, $z = 2.61$

A single, strong emission line is seen in the spectrum. Owing to a lack of any other strong emission lines, we identify this line as Ly α at a redshift of $z = 2.61$. The line also appears to be slightly asymmetric, strengthening the case for Ly α . If the line would be [O II] or [O III], we would expect to see [O III] or H α , respectively, in the redder parts of the spectrum. There is no clear emission line in the redder parts, and most of the expected UV lines at the Ly α redshift lines fall in regions with strong sky lines. Gaussian fit to the emission line gives a line flux of $20.4 \pm 1.6 \times 10^{-17}$ erg s⁻¹ cm⁻² and a relatively narrow FWHM of 810 ± 30 km s⁻¹.

(ix) USS206, $z = 4.01$

A single, bright and asymmetric emission line with two peaks is observed. No other emission lines or continuum is seen in the spectrum. Owing to the double-peaked morphology, H I absorption signatures in the profile and a red ‘wing’, we identify the line as Ly α at $z = 4.01$. The measured line flux is $19.5 \pm 3.1 \times 10^{-17}$ erg s⁻¹ cm⁻², with a broad FWHM of 1110 ± 200 km s⁻¹. If this line were the [O II] doublet, we would expect to see [O III] in the spectrum at ~ 8200 Å. [O III] is ruled out based on the line profile as well as the absence of the expected H α line in the spectrum in this case.

(x) USS172, $z = 4.11$

Table 3. Details of detected emission lines, radio flux densities and luminosities at 150 MHz, and radio sizes of galaxies in this paper. We mark sources with single emission lines that may have uncertain identifications with a (*). The source IDs, radio flux densities, and sizes are taken from Saxena et al. (2018a), which also gives the RA and Dec information for each source. We mention the telescope that was used to observe each source in the Tel. column. ‘W’ stands for WHT, ‘G’ for Gemini North, and ‘H’ for HET.

Source ID	z	Tel.	Line	Flux (10^{-17} erg s $^{-1}$ cm $^{-2}$)	Luminosity (10^{41} erg s $^{-1}$)	FWHM (km s $^{-1}$)	$S_{150 \text{ MHz}}$ (mJy)	$\log L_{150 \text{ MHz}}$ (W Hz $^{-1}$)	LAS_{rad} (kpc)
USS7	0.53	W	[O II] $\lambda 3727$	4.5 ± 3.3	0.5 ± 0.3	970 ± 60	122 ± 24	26.2 ± 0.7	21.6 ± 0.6
–	–	–	[O III] $\lambda 5007$	8.3 ± 2.1	0.9 ± 0.2	460 ± 30	–	–	–
USS43	1.46	W	Mg II $\lambda 2800$	5.0 ± 3.4	6.9 ± 4.7	640 ± 50	200 ± 40	27.6 ± 0.7	5.8 ± 0.8
–	–	–	[O II] $\lambda 3727$	40.0 ± 3.0	55.0 ± 4.1	1380 ± 80	–	–	–
USS483(*)	1.49	G	[O II] $\lambda 3727$	36.6 ± 1.7	13.0 ± 0.6	1050 ± 60	89 ± 16	26.4 ± 0.7	18.8 ± 0.8
USS182	2.02	H	Ly α	32.6 ± 5.0	96.5 ± 14.8	1420 ± 60	146 ± 30	27.8 ± 0.7	24.2 ± 0.8
USS18	2.06	H	Ly α	77.9 ± 4.3	242.0 ± 13.4	950 ± 60	131 ± 26	27.8 ± 0.7	9.5 ± 0.8
USS320	2.20	W, H, G	Ly α	8.7 ± 3.8	31.7 ± 13.9	1300 ± 100	86 ± 17	27.9 ± 0.7	19.6 ± 0.8
–	–	–	Mg II $\lambda 2800$	3.2 ± 1.1	12.1 ± 4.1	320 ± 30	–	–	–
USS268	2.35	W, H	Ly α	22.8 ± 2.8	97.6 ± 12.0	750 ± 50	92 ± 18	27.8 ± 0.7	12.6 ± 0.8
–	–	–	Mg II $\lambda 2800$	5.2 ± 3.0	23.2 ± 13.4	320 ± 30	–	–	–
USS98	2.61	W, G	Ly α	20.4 ± 1.6	112.7 ± 8.8	810 ± 30	178 ± 36	28.3 ± 0.7	56.9 ± 0.8
USS206	4.01	W	Ly α	19.5 ± 3.1	301.7 ± 50.0	1110 ± 200	102 ± 20	28.6 ± 0.7	58.7 ± 0.8
USS172(*)	4.11	H	Ly α	5.2 ± 0.7	85.3 ± 11.5	570 ± 30	124 ± 25	28.6 ± 0.7	65.9 ± 0.7
USS188	4.57	H	Ly α	6.8 ± 0.7	143.3 ± 14.8	520 ± 20	166 ± 33	28.8 ± 0.7	26.2 ± 0.7
USS46	4.86	H	Ly α	5.1 ± 0.8	124.2 ± 20.0	500 ± 30	87 ± 17	28.7 ± 0.7	6.2 ± 0.6
USS202	5.72	G	Ly α	1.6 ± 0.6	57.0 ± 21.4	370 ± 30	170 ± 34	29.1 ± 0.7	6.3 ± 0.6




Figure 3. Top: VLA image of USS483 at 1.4 GHz, with a beam size of ~ 1.3 arcsec. Also shown is the position of the GMOS long-slit, aligned along the radio axis. Bottom: The 2D line emission that we identify as [O II] at $z = 1.49$. The line emission is spatially extended along the radio axis, with emission extending all the way to ~ 15 arcsec away from the central host galaxy. This behaviour is a clear sign of interaction between the radio jet and the surrounding gas.

A single, narrow emission line is detected in the spectrum. No other emission lines are seen in either the blue or the red channels of the spectrograph. We tentatively identify the emission line as Ly α at $z = 4.11$. The line is faint, with an integrated flux of $5.2 \pm 3.0 \times 10^{-17}$ erg s $^{-1}$ cm $^{-2}$ and has a narrow FWHM of 570 ± 30 km s $^{-1}$. We cannot definitively identify this line as Ly α based on the line profile, as the line is relatively symmetric compared to the other line profiles presented in this study and the spectrum is noisier than others presented in this study.

However, this source has NIR photometry from our LBT observations, and the host galaxy is detected with $J = 22.8 \pm 0.1$ AB and $K = 22.7 \pm 0.2$ AB, which are in line with expectations based on the K - z relation for radio galaxies. If the line were to be [O II] or [O III], a redshift of 0.68 or 0.24 would mean that the K -band magnitude of the host galaxy is too faint, making it a strong outlier on the K - z distribution for radio galaxies (see Willott et al. 2003, for example). The NIR colours ($J - K = 0.1$) are also indicative of a high-redshift nature, as a lower redshift radio galaxy would probe the rest-frame NIR where redder colours are expected (see Yang, Chen & Huang 2015, for example).

(xi) USS188, $z = 4.57$

A single emission line is seen in the spectrum. The line is slightly asymmetric, and owing to the absence of any other strong lines in the spectrum, we identify it as Ly α at $z = 4.57$. If this line were [O II], we would expect to see [O III] in the spectrum in the red channel of LRS2 of the HET, and in case the line were [O III], we would expect to see H α . The integrated line flux is $6.8 \pm 0.7 \times 10^{-17}$ erg s $^{-1}$ cm $^{-2}$ with a relatively narrow FWHM of 520 ± 20 km s $^{-1}$. NIR photometry of the host galaxy shows that it is undetected down to a 3σ depth of $J > 23.4$ AB, which is consistent with a high-redshift nature. However, the host galaxy is detected with $K = 21.6 \pm 0.2$ AB, making it relatively bright.

This bright K -band magnitude can be explained through expected contamination from [O II] $\lambda 3727$, which would be redshifted into the K band. Contribution from strong emission lines to broad-band NIR magnitudes is expected from radio galaxies at certain

redshifts (see Miley et al. 2006, for example). It is possible that the strong $J - K$ colour is due to a very dusty host galaxy, which would imply that detection of rest-frame UV/optical emission lines would be highly unlikely in the spectrum (see Reuland et al. 2003, for example). Since an emission line is seen in the spectrum, contamination by a strong emission line in the K band is the likely explanation.

(xii) USS46, $z = 4.86$

We detect a single, asymmetric emission line in the spectrum. No other emission lines or continuum is detected in the wavelength ranges covered. The asymmetric line profile, showing absorption in the blue part of the line and bumps in the red part, characteristic of $\text{Ly } \alpha$ at high redshifts, led us to identify this line as $\text{Ly } \alpha$ at $z = 4.86$. The line has an integrated flux of $5.1 \pm 0.8 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ and a narrow FWHM of $500 \pm 30 \text{ km s}^{-1}$. This makes USS43 one of the highest redshift radio galaxies currently known. Again, $[\text{O II}]$ or $[\text{O III}]$ are ruled out due to the absence of $[\text{O III}]$ or $\text{H } \alpha$, respectively, in the redder channel of the spectrograph.

(xiii) USS202, $z = 5.72$

A single emission line, characteristic of $\text{Ly } \alpha$ at high redshift, is detected in the spectrum of this object, giving a redshift of $z = 5.72$. This makes it the highest redshift radio galaxy observed to date. A detailed analysis of the emission line of this particular source is presented separately in Saxena et al. (2018b).

2.3.4 Undetected sources

There remain 10 USS sources that were targeted for spectroscopy and no emission lines were detected. From these, two sources have continuum detections but no emission lines, implying a possible low redshift. The rest of the sources could either lie in the redshift range where the bright optical emission lines are hard to observe from ground-based telescopes ($1.5 < z < 2$), have passive galaxy type spectra lacking in strong emission lines, have very broad lines that were missed by our relatively shallow spectroscopy, or be heavily obscured. The non-detections could also indicate that these sources are either (optically) faint, low-mass radio galaxies, or lie at very high redshifts ($z > 6.5$; De Breuck et al. 2001).

When following up samples of USS-selected radio sources, there are often sources that do not show UV/optical emission lines but are detected in near-IR broad-band imaging. This is usually a sign of heavy obscuration by dust, and Reuland et al. (2003) showed that following such sources up at submillimetre and millimetre wavelengths is much more efficient in detecting the host galaxy. Using such observations to constrain the SEDs, an accurate photometric redshift can be obtained along with good estimates on the physical properties of the host galaxies. Potential follow-up of the spectroscopically undetected sources at millimetre wavelengths could help identify their host galaxies and reveal their nature.

2.4 Properties of the faint radio galaxy sample

2.4.1 Redshift distribution

In this section, we look at the redshift distribution of the spectroscopically confirmed radio sources. Out of a total of 32 sources that were initially compiled by Saxena et al. (2018a), we spectroscopically confirmed redshifts for 13 of them. Fig. 4 shows the distribution of these redshifts, along with photometric redshifts for three additional objects in the sample reported by Saxena et al. (2018a), which pushes the total number of sources with redshifts to 16. The total range of spectroscopic redshifts is $0.52 < z < 5.72$. This includes the recent




Figure 4. Stacked histogram of spectroscopic redshifts (blue) and photometric redshifts (orange) for radio sources in the Saxena et al. (2018a) sample. The total height of the bar shows the total number of objects with either spectroscopic or photometric redshifts in each bin. The distribution shows that the USS selection is effective at picking out high-redshift radio galaxies from a large-area radio survey.

discovery of the highest redshift USS radio source at $z = 5.72$, which was reported separately in Saxena et al. (2018b). The detection of a fair number of $z > 2$ radio galaxies, through USS selection combined with optical faintness and compact radio sizes, demonstrates the effectiveness of these selection criteria and holds promise for similar studies in the future with even deeper radio surveys, such as the LOFAR Two-meter Sky Survey (Shimwell et al. 2017, 2019).

2.4.2 Velocity dispersion

For radio sources spectroscopically confirmed in this study, through the $[\text{O II}]$, $[\text{O III}]$, and Mg II lines at low redshifts and the $\text{Ly } \alpha$ line at high redshifts ($z > 2$), the FWHM of the emission lines is in the range $320\text{--}1420 \text{ km s}^{-1}$. This range is lower than what is typically observed for much brighter radio galaxies across all redshifts (see De Breuck et al. 2000a, for example), and in the following sections we compare the line widths with radio galaxies in the literature.

2.4.3 Stellar masses

We now estimate stellar masses for 8 (out of the 13 spectroscopically confirmed) new radio galaxies that were observed in the K band. To do this, we first model the expected K -band magnitudes for radio galaxies with different stellar masses across redshifts using the PYTHON package called SMPY (Duncan & Conselice 2015). SMPY is a stellar population modelling code designed for building composite stellar populations in an easy and flexible manner and allows for synthetic photometry to be produced for single or large suites of stellar population models. To build stellar populations, we use the Bruzual & Charlot (2003) model with a Chabrier (2003) initial mass function (IMF). We assume solar metallicity, similar to Willott et al. (2003), a formation redshift of $z_f = 25$, and a maximally old stellar population that has been forming stars at a constant rate. We follow the Calzetti et al. (2000) law for dust attenuation, using $A_V = 0.5$, in line with observations and predictions from McLure et al. (2018) and Cullen et al. (2018) at the redshifts and mass bins that are typical of radio galaxy hosts ($M_* > 10^{10.2} M_\odot$). The synthetic photometry produced for stellar masses in range $10.6 <$

Table 4. NIR photometry for HzRGs and stellar mass measurements for radio galaxies in our sample.

Source ID	z	J (AB)	K (AB)	$\log M_*$ ($M_\odot$)
USS43	1.46	—	21.6 ± 0.2	10.6 ± 0.4
USS182	2.02	—	21.8 ± 0.2	11.0 ± 0.4
USS320	2.20	—	21.7 ± 0.3	11.1 ± 0.4
USS268	2.35	23.6 ± 0.1	21.9 ± 0.3	11.1 ± 0.5
USS172	4.11	22.8 ± 0.1	22.7 ± 0.2	11.2 ± 0.5
USS188	4.57	>23.4	21.6 ± 0.2^a	11.5 ± 0.5
USS202	5.72	>25.3	>24.2	<10.5

Note. ^aExpected contamination from [O II] $\lambda 3727$ emission line in the K band.

$\log M_* < 12.0$ is then convolved with the K -band filter to calculate apparent K -band magnitudes over a redshift range $0 < z < 7$. The individual stellar mass is determined by finding the best-fitting stellar mass track to the K -band magnitude and redshift using chi-squared minimization.

The NIR photometry for faint radio galaxies and stellar masses inferred from the models are shown in Table 4. The fainter radio galaxies have lower stellar masses on average when compared to the bright radio galaxies known in the literature. The $10^{12} M_\odot$ stellar mass limit predicted for radio galaxies by Rocca-Volmerange et al. (2004) stands even for our sample – there is no radio source with an inferred stellar mass higher than $10^{12} M_\odot$ in our sample. Overall, we find that fainter radio galaxies occupy the massive end of the stellar mass function at all redshifts, in line with previous observations that showed that radio galaxies are hosted by massive galaxies at every epoch (Rocca-Volmerange et al. 2004). The highest redshift radio galaxy, TGSS J1530+1049 at $z = 5.72$, which was part of our sample and was presented in Saxena et al. (2018b), is one of the lowest stellar mass radio galaxy from our sample.

We note that the overall lower stellar masses of galaxies in our sample are obtained under the assumptions that go into the stellar population modelling mentioned earlier. We note that there are several key parameters that contribute towards the uncertainty on the determined stellar mass. Multiband photometry is generally required to obtain accurate stellar masses, by constraining the SED of hosts of radio galaxies, taking into account contributions from the central AGN, stars, and dust (see Seymour et al. 2007; Overzier et al. 2008, for example). Even then, assumptions regarding the IMF, formation redshift, star formation history, and the type of underlying stellar populations all contribute towards the uncertainties in stellar mass.

Since our estimates are based on only one or two broad-band photometric points, and the goal of our stellar mass measurement is to get a very simple idea based on K -band photometry, taking into account all the uncertainties is beyond the scope of this current analysis. However, since our models are able to account for the scatter in the K - z relation for known radio galaxies without requiring extremely large or extremely small stellar masses, the inferred stellar masses for galaxies in our sample seem to be in line with expectations. Follow-up broad-band photometry across optical to mid-infrared wavelengths, capturing light from the different sources that contribute to the SED, is required to put robust constraints on the stellar mass of these radio galaxies.

3 COMPARISON WITH LITERATURE HZRG SAMPLE

In this section, we compile a sample of known high-redshift radio galaxies ($z \geq 2$) from the literature to compare the sample of newly discovered faint HzRGs against. We begin by briefly describing the

parent radio samples from which the literature $z > 2$ radio galaxies were selected. We particularly highlight the flux limits of each of these radio samples and note that a large majority of these radio samples have much higher flux density limits than the Saxena et al. (2018a) sample used in this study. We point the readers to the relevant references mentioned in the sample description for detailed information about the sources.

3.1 Literature samples

3.1.1 Röttgering et al. (1997)

This sample contains spectroscopic follow-up of the Leiden USS sample (Röttgering et al. 1994) and the 4C USS sample Tielens et al. (1979). Röttgering et al. (1997) sample contains spectra for 64 radio galaxies, with 29 of them being at $z > 2$. Other Ly α properties from this sample are presented in van Ojik et al. (1997). The median radio flux density of these sources is in the range 1–2 Jy.

3.1.2 De Breuck et al. (2001)

This spectroscopic sample of radio galaxies follows up the USS radio sources that were first presented in De Breuck et al. (2000a). The De Breuck et al. (2001) sample contains spectra for 69 radio galaxies from the De Breuck et al. (2000a) radio sample and observations of a few other previously known HzRGs, with 22 of them confirmed at $z > 2$. The limiting radio flux density for various subsamples observed in this study ranges from 250 to 1000 mJy.

3.1.3 Jarvis et al. (2001a)

This sample contains spectra for radio sources selected from the 6C* sample (Blundell et al. 1998). The Jarvis et al. (2001a) sample has a total of 29 spectra, 12 of which lie at $z > 2$. The flux density at 151 MHz of sources in this sample is in the range 0.96–2 Jy.

3.1.4 Willott et al. (2002)

This sample contains optical spectroscopy for 49 radio sources selected from the 7C Redshift Survey, which is a spectroscopic survey of the regions covered in the 5C and 7C radio surveys. The Willott et al. (2002) sample contains five radio galaxies at $z > 2$. The radio flux density probed by this sample is $S_{151} > 0.5$ Jy.

3.1.5 Bornacini et al. (2007)

This sample contains spectra for 14 USS sources selected from the Westerbork in the Southern Hemisphere (WISH) catalogue (De Breuck et al. 2002) and subsamples from the De Breuck et al. (2000a) USS catalogue. 6 out of 14 sources observed by Bornacini et al. (2007) lie at $z > 2$. The limiting flux density of the WISH sample is $S_{352} \sim 18$ mJy.

3.1.6 Bryant et al. (2009)

This sample contains redshifts for 36 galaxies, out of a total of 234 USS radio sources shortlisted from the Revised Molonglo Reference Catalogue, Sydney University Molonglo Sky Survey (MRCR-SUMMS sample) and the NRAO VLA Sky Survey (NVSS), presented in Broderick et al. (2007). There are 13 radio sources at $z > 2$ from the Bryant et al. (2009) sample, 11 of which have Ly α line




Figure 5. The distribution of radio luminosities and redshifts of the faint HzRGs (gold stars) and the literature galaxies (coloured symbols), which have been scaled to an observing frequency of 150 MHz. The faint HzRGs are almost an order of magnitude fainter at the highest redshifts, opening up a new parameter space for the discovery and study of radio galaxies in the early Universe.

measurements. The Broderick et al. (2007) parent radio catalogue contains sources with $S_{408} \geq 200$ mJy.

3.1.7 Radio luminosities

The extended HzRG sample consists of a total of 86 radio galaxies at $z \geq 2$. All these radio galaxies have $\text{Ly}\alpha$ line measurements. We calculate the radio luminosities, radio sizes, and $\text{Ly}\alpha$ luminosities for the literature HzRGs using the updated cosmology used in this study. Specifically for the radio luminosities at a frequency of 150 MHz, we use the published flux density at 150 MHz wherever available and when needed, we scale their flux density to 150 MHz using the published spectral index. For the newly spectroscopically confirmed radio galaxies reported in this paper, we use the radio measurements from Saxena et al. (2018a) to calculate radio luminosities.

The distribution of radio luminosities as a function of redshift is shown in Fig. 5. Our sample of newly confirmed faint HzRGs is marked with gold stars (and in all the following sections). It is immediately clear that our sample probes radio luminosities almost and order of magnitude fainter at the highest redshifts. This includes the discovery of the highest redshift radio galaxy known till date, TGSS J1530+1049 (Saxena et al. 2018b). We also show the faint radio galaxy detected by Jarvis et al. (2009), which was detected in a deep field and not from an all-sky survey, as an example of the kind of sensitivities that future radio surveys would be able to achieve to probe even fainter radio galaxies at high redshifts.

3.2 Comparison of emission line and radio properties

In this section, we will explore how the emission line and radio properties of faint HzRGs compare with those of the literature HzRG sample.

3.2.1 $\text{Ly}\alpha$ luminosities and FWHM

We begin our analysis by first showing the distribution of $\text{Ly}\alpha$ line luminosities and FWHM values for faint HzRGs (gold stars) and literature HzRGs (black circles) across all redshifts in Fig. 6. We note that the faint HzRGs in our sample show a systematically lower $\text{Ly}\alpha$

luminosity and FWHM compared to other HzRGs in the literature. The faint HzRG sample has a median $\text{Ly}\alpha$ luminosity of $\log L_{\text{Ly}\alpha} = 43.0 \text{ erg s}^{-1}$, with 1σ standard deviation of 0.3 dex. The literature HzRG sample has a median $\text{Ly}\alpha$ luminosity of $\log L_{\text{Ly}\alpha} = 43.4 \text{ erg s}^{-1}$ but a much larger 1σ standard deviation of 0.7 dex. The difference in median FWHM values between the two samples is large, with faint HzRGs having median and standard deviation of 780 ± 340 (1σ) km s^{-1} , and the literature HzRGs having median and standard deviation of $1420 \pm 910 \text{ km s}^{-1}$.

In Fig. 7, we plot the $\text{Ly}\alpha$ luminosities with redshift for both faint and literature samples. The higher dispersion in the literature sample is seen across all redshifts, with a handful of sources at $z \sim 3$ – 3.5 showing significant deviations (2.5σ – 3σ) from the median, such as WN J1053+5424 at $z = 3.1$ exhibiting the weakest $\text{Ly}\alpha$ luminosity of the sample ($\log L_{\text{Ly}\alpha} = 41.7 \text{ erg s}^{-1}$) and TN J0205+2242 at $z = 3.5$ having the strongest luminosity ($\log L_{\text{Ly}\alpha} = 45.9 \text{ erg s}^{-1}$). The faint HzRGs have comparable $\text{Ly}\alpha$ luminosities across redshifts. Overall, the luminosities (both faint and literature) do not seem to evolve much with redshift, with the Pearson’s correlation coefficient $r = -0.02$ suggesting little to no evolution. Previously, Zirm et al. (2009) had reported the detection of lower luminosity $\text{Ly}\alpha$ emission from a sample of $z \sim 1$ radio galaxies that exhibited strong nebular emission lines, suggesting that the $\text{Ly}\alpha$ luminosities of extended haloes around radio galaxies may undergo strong redshift evolution. Although our samples of both faint and literature radio galaxies do not exhibit very strong redshift evolution of $\text{Ly}\alpha$ luminosity, expanding the sample of faint radio galaxies at high redshifts may hold some clues about any such behaviour.

Next, we plot the $\text{Ly}\alpha$ line widths (FWHM) as a function of redshift for both faint and literature HzRG samples in Fig. 8. We find a negative correlation between FWHM and redshift for both samples and the anticorrelation is stronger in the faint sample ($r = -0.77$, compared to $r = -0.20$ for the literature sample). From the literature sample, there are 3 radio galaxies at $z \sim 2$ – 3 that are 3σ outliers from the median value ($\text{FWHM} > 4141 \text{ Å}$), with the radio galaxy 1113–178 from Röttgering et al. (1997) showing the broadest $\text{Ly}\alpha$ profile with $\text{FWHM} = 5000 \text{ km s}^{-1}$.

3.2.2 Radio luminosity versus $\text{Ly}\alpha$ luminosity

We now show $\text{Ly}\alpha$ line luminosities as a function of radio luminosity at 150 MHz for the faint and literature samples in Fig. 9. We see a weak positive correlation in the literature sample, with $r = 0.17$, which is weaker than what Jarvis et al. (2001a) observed for their sample of radio galaxies. Conversely, in the faint HzRG sample we see a very weak correlation, with $r = 0.04$, which is compatible with no correlation. Both these r -values, that show little to no correlation between the two quantities, indicate that the radio luminosity (across several orders of magnitude) of a radio galaxy does not have much influence on its $\text{Ly}\alpha$ luminosity.

3.2.3 Linear sizes

In this section, we look at the linear size distribution of radio galaxies in both samples. For faint HzRGs presented in this paper, we use the angular sizes reported by Saxena et al. (2018a). Linear sizes of literature HzRGs were calculated using the updated cosmology used in this study and their published angular sizes. The median size of our faint radio sources is 22 kpc and the median size of the literature HzRGs is 42 kpc. We highlight that there was angular size restriction ($< 10 \text{ arcsec}$) implemented by Saxena et al. (2018b) while selecting




Figure 6. Distribution of $\text{Ly}\alpha$ luminosities and FWHM for faint (gold stars) and literature (black circles) radio galaxies. Also shown are the histograms of both these samples. Clearly, the faint HzRGs have lower $\text{Ly}\alpha$ luminosities and FWHM. In the text, we give the median and standard deviations of $\text{Ly}\alpha$ luminosities and FWHM in the two samples. There is no correlation observed between the $\text{Ly}\alpha$ luminosities and FWHM for either of the two samples.




Figure 7. $\text{Ly}\alpha$ line luminosities as a function of redshift for the faint and literature HzRG samples. Across all redshifts, the faint HzRGs show lower luminosity values. We do not see any evolution in the line luminosity with redshift for either of the two HzRG samples.

faint HzRG candidates, which would preferentially select compact radio sources across all redshifts.

We show the distribution of radio luminosities at 150 MHz as a function of linear size for both the faint and literature HzRGs in Fig. 10. These so-called ‘ $P - D$ ’ diagrams have long been used to characterize the growth and evolution of powerful radio sources across all redshifts (Kaiser, Dennett-Thorpe & Alexander 1997; Blundell et al. 1999; Saxena et al. 2017). The $P - D$ tracks




Figure 8. The $\text{Ly}\alpha$ FWHM as a function of redshift for the faint and literature HzRGs. Across all redshifts, faint HzRGs exhibit lower FWHM values when compared to the literature HzRGs. We see a decrease in FWHM with redshift in both samples, but the anticorrelation is stronger for faint HzRGs (orange dashed line) than the bright ones (black dashed line).

essentially trace the evolutionary path that a radio source takes through its lifetime. For a given jet power, radio sources start out at the left side of the $P - D$ diagram. As radio sources grow, they lose energy, initially through adiabatic losses and at larger sizes due to inverse Compton (IC) losses because of interactions with the cosmic microwave background radiation (Kaiser & Best 2007). Overall, this results in the radio sources experiencing decreasing




Figure 9. Ly α luminosities as a function of radio luminosity at 150 MHz for faint and literature HzRGs. There is a positive correlation observed between the two quantities for literature HzRGs; however, it is weaker than what some previous studies have found. However, we do not see any correlation between the two for faint HzRGs.




Figure 10. The distribution of radio luminosities at 150 MHz and linear sizes of sources in the faint and literature samples. The $P - D$ diagram tracks the evolution of radio sources through their lifetime. Clearly, the most luminous sources are able to grow out to larger sizes as their intrinsically more powerful jets work against the energy loss mechanisms that begin to dominate later in the radio source life. The faint HzRGs preferentially lie towards the low power, small size part of the plot, indicating that they must be young.

radio luminosities with increasing linear sizes, moving them towards the bottom-right side of the $P - D$ diagram. The $P - D$ tracks are influenced by magnetic field strength, redshift, and the radio source environment.

As seen in Fig. 10, the faint HzRGs lie in the part of the $P - D$ diagram corresponding to weaker and younger radio sources. We also note that the largest radio sources are also some of the most luminous ones and larger radio sources are expected to be older than the compact ones. Radio sources with intrinsically weaker jets are unable to grow out to large sizes as they are unable to overcome the increasing energy losses with increase in linear size, and are eventually extinguished before ever achieving large sizes (Saxena et al. 2017). This is evident from the lack of sources in the bottom-right part of the $P - D$ diagram. We discuss the redshift evolution of linear sizes of both the faint and literature samples in Section 4.4.




Figure 11. Low-frequency radio spectral index as a function of redshift for both faint and literature HzRGs. We observe an anticorrelation between z and α for the literature sample, which has been used successfully to select promising HzRG candidates from large radio surveys. However, the faint HzRGs actually show little to no correlation between the two quantities. Since most of the HzRGs presented in this study were initially selected because of their spectra, the observed correlations are likely to contain several biases.

3.2.4 Radio spectral index versus redshift

We now explore the observed correlation between the low-frequency radio spectral index (α) and redshift (z) in our extended HzRG sample in Fig. 11. The $z - \alpha$ correlation for radio galaxies has often been used to identify HzRG candidates in large-area radio surveys (see Röttgering et al. 1997, for example).

We observe an anticorrelation between redshift and spectral index in the radio galaxies from the literature, with $r = -0.50$. However, it is important to note that most of the radio galaxies that are part of the extended sample were initially selected because of their USS at low radio frequencies ($\alpha < -1$), which bias the distribution and lead to the recovered correlation. From our sample of faint galaxies, which were also selected for having USSs, we do not see a strong anticorrelation between spectral index and redshift. In fact, the faint radio galaxies exhibit a positive correlation between α and z , in contrast to what is seen for literature HzRGs. Further, we do not find extremely steep spectral indices at the highest redshifts, even though the best fit from the literature would suggest so.

The origin of the observed $z - \alpha$ correlation was investigated by Morabito & Harwood (2018), where they found that a combination of IC losses at high redshifts and selection effects was able to explain the observed $z - \alpha$ correlation in radio galaxies. IC losses affect the high-frequency parts of the radio SED, which results in an overall steeper spectral index when measured at low frequencies. IC losses are also expected to affect the linear sizes of radio sources. However, we do not find any strong correlation between the spectral index and linear sizes of radio sources in either of the two samples studied.

4 DISCUSSION

4.1 Ly α emission in faint HzRGs is powered by a weak AGN

As shown in Fig. 6, our newly discovered faint radio galaxies have lower Ly α luminosities and FWHM values across all redshifts compared to the bright HzRGs known in the literature. The dispersion in their Ly α luminosities is also lower. This is interesting because the distribution of Ly α luminosities and FWHM for brighter HzRGs

shows a much larger variety (3σ dispersion of $\log L_{\text{Ly}\alpha} = 2.1$ erg s^{-1} , as opposed to 0.6 erg s^{-1} for the faint radio galaxies, as shown in Section 3.2.1). Historically, the large luminosities and equivalent widths of $\text{Ly}\alpha$ emission from bright HzRGs have been explained by anisotropic photoionization from an obscured AGN in combination with jet-induced shock ionization or contribution from hot stars (Chambers, Miley & van Breugel 1990; Baum et al. 1992; McCarthy 1993; van Ojik et al. 1996, 1997; De Breuck et al. 2000b). Additionally, physical phenomena such as launching of winds and outflows from the central AGN (see Nesvadba et al. 2008, for example), bursts of star formation and radio AGN activity driven by recent mergers (see Villar-Martín et al. 2007b, for example), and gas infall on to the host galaxy (see Villar-Martín et al. 2007a, for example) are sometimes required to explain the UV line ratios and large velocity dispersions at high redshifts.

The systematically lower $\text{Ly}\alpha$ luminosities in faint HzRGs indicate that the line emission must be ionized by a weaker AGN, or by modest star-burst activity. To accurately disentangle the relative contributions from an AGN and star formation, diagnostics based on UV emission line ratios are required. Several studies in the literature have showed that UV line ratios can be used to distinguish between shocks or AGN photoionization being the dominant mechanisms (Villar-Martín et al. 1997; Allen, Dopita & Tsvetanov 1998; Best, Röttgering & Longair 2000; De Breuck et al. 2000b). It has also been shown that UV line ratios can be used to distinguish between star formation or AGN activity being the dominant ionising mechanism powering the lines (Feltre, Charlot & Gutkin 2016). For the redshifts of interest presented in this study, follow-up NIR spectroscopy in the future offers the best way to detect rest-frame UV emission lines and test various scenarios of ionization. In the following sections, however, we try to set some qualitative constraints on the dominant ionizing mechanisms of faint HzRGs.

4.2 No direct evidence of strong interaction between radio jet and line-emitting gas

Another indication of the absence of shocks is the lack of evidence for strong jet–gas interaction – we do not see an anticorrelation between $\text{Ly}\alpha$ FWHM and linear size, which has previously been interpreted as a signature of jet–gas interaction in radio galaxies (van Ojik et al. 1997; Baum & McCarthy 2000; Best et al. 2000). The origin of this anticorrelation has been attributed to jets in smaller radio sources still being in the process of propagating through the relatively dense ISM of their host galaxies, which leads to transfer of kinetic energy from the jets to the line-emitting gas, resulting in larger line widths (see van Ojik et al. 1997, for example). For larger radio sources, when the jet has passed well beyond the emission-line regions, the emission line widths are narrower indicating that the gas has more relaxed and settled kinematics.

In fact, when combining the faint and literature HzRG samples, it is clear that even though some of the largest radio sources in our sample show narrower $\text{Ly}\alpha$ line widths, we cannot conclusively infer that narrow line widths are a consequence of less jet–gas interaction, as shown in Fig. 12 in which the faint HzRGs are marked with gold stars and literature HzRGs with black circles. The median linear sizes in the combined sample were calculated in bins of $\text{Ly}\alpha$ FWHM. These FWHM bins are 0–1000, 1000–2000, 2000–3000, and >3000 km s^{-1} . Therefore, from our data we cannot conclusively pinpoint the dominant impact of jet–gas interactions in smaller radio sources. However, the added complexity of probing radio sources across a broad range of radio and $\text{Ly}\alpha$ luminosities as well as redshifts in our




Figure 12. The linear sizes of faint HzRGs (gold stars) are not seen to anticorrelate with $\text{Ly}\alpha$ FWHM, which is indicative of absence of strong jet–gas interaction. This implies that shocks do not play a strong role in powering the $\text{Ly}\alpha$ emission in faint HzRGs. The anticorrelation in the combined sample of faint and literature HzRGs, calculated by binning in FWHM, is also very weak (black squares), although we see that the largest literature HzRGs often also exhibit narrower FWHM.

sample may dilute out any strong effects that may be seen in subsets of relatively similar radio galaxies.

4.3 Faint HzRGs have lower stellar masses

As shown earlier, the apparent K -band magnitudes for a subset of faint radio galaxies presented in this paper were lower than what has been observed for more powerful radio sources across all redshifts. Using simple stellar population modelling, we find that the stellar masses inferred for faint HzRGs are also lower than what has been typically determined for bright HzRGs (see Rocca-Volmerange et al. 2004, for example). This is shown in Fig. 13, where we show apparent magnitudes for our faint HzRGs along with those from HzRGs in the literature.

Also shown are predictions for apparent K magnitude from our stellar population synthesis modelling. Our faint HzRGs lie closer to the lower mass tracks, with the exception of USS188 with a stellar mass of $M_* = 10^{11.7 \pm 0.7} M_\odot$. However, contamination from the $[\text{O II}]\lambda 3727$ emission line is expected in the K band and it is therefore likely that the stellar mass for this galaxy is being overestimated. The stellar mass estimates presented in this paper are based on only the K -band magnitudes and photometry in additional bands and more detailed stellar population modelling is required to accurately measure stellar masses for our sample of faint HzRGs.

4.4 The nature of faint HzRG hosts

As shown earlier, the faint HzRGs reported in this paper overall show weaker $\text{Ly}\alpha$ luminosities and narrower line widths in comparison to the brighter HzRGs. These fainter radio galaxies are also less massive than the brighter ones. Questions naturally arise about the nature of faint radio galaxies at high redshifts. In this section, we attempt to provide some qualitative constraints on this.

To better understand the nature of fainter radio galaxies, we compare their $\text{Ly}\alpha$ line luminosities and widths with bright, ‘non-radio’ $\text{Ly}\alpha$ emitters (LAEs) at comparable redshifts from the literature. The comparison sample is constructed mainly from LAEs that have deep spectroscopy over a wide range of luminosities, in the redshift range




Figure 13. Apparent K -band magnitudes and redshifts for faint HzRGs (gold stars) and those from the literature (black dots). The solid lines indicate photometry obtained using simple stellar population modelling as described in Section 2.4.3. The solid lines show stellar mass tracks in the range $\log M_* = 10.2$ – $12.0 M_\odot$, in increments of $\log M_* = 0.2 M_\odot$. Our fainter radio galaxies have lower stellar masses when compared to the previously known, more powerful radio galaxies across all redshifts. We note that TGSS J1530+1049 at $z = 5.72$ has only a stellar mass limit, as it was undetected in our NIR imaging campaign (Saxena et al. 2018b). The galaxy with the brightest K mag in our sample (USS188 at $z = 4.57$) has expected contamination from the [O II] emission line in the broad-band, which probably leads to an overestimated stellar mass using simple models.

covered by the faint HzRGs reported in this paper. The samples we consider by no means represent the complete LAE population, but offer good statistics for comparison with our radio galaxies.

In this analysis, we include the 21 LAEs observed by Sobral et al. (2018) in the redshift range $z \sim 2$ – 3 , which were initially selected from narrow-band surveys and then followed up spectroscopically. Sobral et al. (2018) identify three broad classes of objects in their sample – broad-line AGN, narrow-line AGN, and star-forming galaxies. We also include the sample of 18 bright LAEs in the redshift range $z \sim 3$ – 5 observed by Saito et al. (2008). This sample is divided into low equivalent width ($EW < 200 \text{ \AA}$) and high- EW ($> 200 \text{ \AA}$) sources and both subsamples contain star-forming galaxies. Finally, we include the spectroscopically confirmed brightest LAEs from the SILVERRUSH survey (Ouchi et al. 2018) presented in Shibuya et al. (2018). These LAEs were narrow-band selected at $z = 5.7$ (comparable with TGSS J1530+1049) and $z = 6.6$, representing some of the brightest LAEs at $z > 5.5$. Since the $\text{Ly}\alpha$ luminosities and line widths are quite similar for the two redshifts probed by the Shibuya et al. (2018) sample, we calculate the median luminosity and line width for each redshift and use that for comparison.

In Fig. 14, we show the $\text{Ly}\alpha$ luminosities and FWHM for the faint HzRGs (gold stars), the literature HzRG sample (black dots), and the LAEs from Sobral et al. (2018) (crosses), Saito et al. (2008) (pluses), and median values from Shibuya et al. (2018) at two redshifts (squares). Clearly, the literature HzRGs show a large scatter in both $\text{Ly}\alpha$ luminosity and FWHM values, whereas the distribution for faint HzRGs has a lower scatter. The faint HzRGs overlap with SF galaxies and narrow-line AGN from Sobral et al. (2018) but not broad-line AGN, whereas a large number of literature radio galaxies occupy the same parameter space as broad-line AGN. Another interesting trend is the tight distribution in the parameter space of star-forming galaxies in both Sobral et al. (2018) and Saito

et al. (2008) samples, and eight of our faint radio galaxies follow this trend all the way into the parameter space occupied by an AGN. Four galaxies from our sample with $\text{FWHM} < 600 \text{ km s}^{-1}$ (below the dashed horizontal line), which all also are at $z > 4$, show a complete overlap with star-forming galaxies from the LAE samples. We now try and qualitatively describe the nature of radio galaxy hosts based on what is known about the ‘non-radio’ LAEs used for comparison.

4.4.1 Faint HzRGs at $z \sim 2$ – 4

We first look at the fainter HzRGs presented in the redshift range $z \sim 2$ – 4 . This range consists of 6 out of the 10 sources at $z > 2$ with $\text{Ly}\alpha$ measurements (lying above the dashed black line). We note that these radio galaxies occupy the regions in the parameter space in Fig. 14 that mark the transition between the star-forming and AGN populations. For these, the Sobral et al. (2018) galaxies offer a good sample for comparison owing to their comparable redshift range. We note that for their bright LAEs, Sobral et al. (2018) use photoionization modelling to show that the star-forming galaxies (green squares) have a starburst age of $\sim 20 \text{ Myr}$, ionization parameters $\log U \sim -3$, and mildly subsolar gas metallicities. For AGNs, they find an ionization parameter $\log U \sim 0.6$ and higher gas metallicities, extending up to $2 Z_\odot$.

Since two or more UV emission lines are essential to accurately determine the dominant source of ionization, and as we only detect a single $\text{Ly}\alpha$ line in a majority of our shallow spectra, we must rely on inferring ionization properties by qualitative comparison alone. Following the measurements of Sobral et al. (2018), we can conclude that faint radio galaxies at $z \sim 2$ – 4 represent a population that is transitioning from being AGN dominated to star formation dominated. This is in contrast to a majority of bright radio galaxies known in the literature at these redshifts that display AGN-like features (see De Breuck et al. 2001, for example). Further, we do not see any source that could be classified as a broad-line AGN. Therefore, faint radio galaxies in this redshift range must either be recent starbursts or narrow-line AGN, or a population transitioning from one to the other. Very little is known about faint radio galaxies at these redshifts and larger samples combined with deeper spectroscopy are essential to understand the nature of such sources.

4.4.2 Faint HzRGs at $z > 4$

We now look at the four radio galaxies (lying below the black dashed line in Fig. 14) that completely overlap with the star-forming LAE population. Perhaps a more direct comparison in terms of redshift for these is offered by the Saito et al. (2008) sources. The low- EW LAEs in the Saito et al. (2008) sample have been identified as moderate starbursts, with star formation rates in the range 3.3 – $13 M_\odot \text{ yr}^{-1}$. The presence of a weak AGN in these sources cannot be completely ruled out. For the high- EW LAEs, Saito et al. (2008) rule out the presence of ‘superwinds’ launched by bursts of star formation or ionization by a central AGN as the dominant ionizing mechanism. Owing to the observed tight correlation between the line luminosities and widths, Saito et al. (2008) conclude that the origin of line emission in high- EW LAEs is most likely due to accretion of cold gas on to the galaxy, placing them in an early phase of their evolution.

The Shibuya et al. (2018) LAEs lie at slightly higher redshifts and have much higher $\text{Ly}\alpha$ luminosities than our radio galaxies. Deep spectroscopy of these LAEs also led to non-detection of other UV lines in their spectra. Shibuya et al. (2018) rule out the presence of broad-line AGN and Population III (very low metallicity stars) as




Figure 14. $\text{Ly}\alpha$ luminosity plotted against line width for faint (gold stars) and literature (black circles) HzRGs. Also shown are ‘non-radio’ $\text{Ly}\alpha$ emitting galaxies (LAEs) from Sobral et al. (2018) in the redshift range $z \sim 2-3$ (crosses), consisting of star-forming galaxies, narrow-line AGN, and broad-line AGN. We also show LAEs observed by Saito et al. (2008) in the redshift range $z \sim 3-5$ (pluses), divided into low EW ($<200 \text{ \AA}$) and high EW ($>250 \text{ \AA}$) and median properties of LAEs from Shibuya et al. (2018) at $z = 5.7$ and 6.6 (squares). The black dashed line at $\text{FWHM} = 600 \text{ km s}^{-1}$ divides the faint radio galaxies at $z \sim 2-4$ (lying above) and $z > 4$ (lying below). We find that $z \sim 2-4$ radio galaxies form a bridge between the star-forming and narrow-line AGN populations, whereas the $z > 4$ radio galaxies overlap fully with the star-forming galaxy population. Our $z > 4$ sources are likely powered by either moderate starburst activity or cold accretion, and must be actively building up their stellar mass.

the drivers of the strong $\text{Ly}\alpha$ emission, and favour intense star-burst activity as the most likely explanation. However, the true nature of their sources remains inconclusive.

In the light of these comparison samples, we may conclude that since our $z > 4$ radio galaxies coincide well with the distribution of both low- and high-EW LAEs from Saito et al. (2008) and are likely to be powered either by ordinary starbursts or by cold accretion. The Saito et al. (2008), Sobral et al. (2018), and Shibuya et al. (2018) sources lying in this parameter space do not favour AGN as the dominant ionizing source. Therefore, we can say with some confidence that faint radio galaxies at $z > 4$ in our sample are starburst dominated, which is different from what has been previously observed in brighter radio galaxies at these redshifts. Further, our radio galaxies show a positive correlation and a tight distribution between $\text{Ly}\alpha$ luminosity and line width, in line with the trend observed by Saito et al. (2008). Therefore, we can also infer that faint HzRGs at $z > 4$, if powered by cold accretion, are in an early stage of their evolution and are actively building up their stellar masses. This is consistent with the lower than average stellar masses calculated from NIR photometry for these radio galaxies.

The suggestion that faint radio galaxies at $z > 4$ are actively building up their stellar mass is not entirely implausible, since radio sources at the highest redshifts are expected to be young (see Blundell et al. 1999, for example). Recent VLBI radio observation of TGSS J1530+1049, a radio galaxy from our sample at $z = 5.72$ (Saxena et al. 2018b), by Gabányi et al. (2018) revealed two distinct lobes

separated by only 0.4 arcsec, confirming that the radio source is young and that the host galaxy must also be in a very early phase of its evolution.

We highlight a caveat of shallow spectroscopy though, which is possibly missing out on sources with broad emission lines. Broad lines may be present in the spectra of sources that we did not detect in our spectroscopy using relatively short exposure times, and observations of these possible broad lines may alter the inferred physical properties of faint radio galaxies that have been reported in this section.

4.4.3 Comparison with $\text{Ly}\alpha$ haloes seen at high redshifts

It has been shown that several radio galaxies at $z > 2$ are associated with extended $\text{Ly}\alpha$ haloes that are kinematically relaxed ($\text{Ly}\alpha$ FWHM $\sim$ few 100 km s^{-1}) and often bright ($\text{Ly}\alpha$ luminosities $\sim 10^{43-44}$; Villar-Martín et al. 2003; Miley et al. 2006). Deep integral-field spectroscopy (IFU), particularly with *MUSE*, has indeed revealed the presence of relaxed $\text{Ly}\alpha$ haloes around both galaxies (Wisotzki et al. 2016) and luminous quasars (Cantalupo et al. 2014; Arrigoni Battaia et al. 2019) at $z > 3$. The existence of such haloes around the faint radio galaxies reported in this study could explain the lower than average $\text{Ly}\alpha$ FWHM that we observe. However, we do not see any signs of extended $\text{Ly}\alpha$ emission in the shallow spectra that we currently have. Indeed, ultra-deep sensitivities are often required to probe extended haloes around high-redshift objects. The $\text{Ly}\alpha$ luminosities seen in the haloes observed in the literature




Figure 15. Linear sizes of the faint HzRGs (gold stars) and literature HzRGs (black circles) with redshift. The median sizes, which are calculated by binning all radio sources in the combined HzRG sample in redshift bins, are shown as large black circles. Also shown are the predictions for redshift evolution of linear sizes due to IC losses, through interaction with the ambient cosmic microwave background radiation, from Saxena et al. (2017), that effectively provide an upper limit on achievable radio size. The median sizes of radio sources evolve strongly with redshift, and are well described by the Saxena et al. (2017) prediction, indicating that the dominant energy loss mechanisms that shape the size distribution of radio sources at high redshifts are IC losses.

also range over several orders of magnitudes and therefore, with the spectra presented in this paper, we cannot confirm the presence of such haloes around faint radio galaxies. Deep follow-up observations, either through IFU spectroscopy or narrow-band imaging at the $\text{Ly}\alpha$ wavelengths of faint HzRGs, could be the key to unveiling any such haloes in which HzRGs may be embedded.

4.5 IC losses dominant at high redshift

Although there is no strong redshift evolution seen in the sizes of the faint HzRGs reported in this paper, we do observe an evolution of radio sizes with redshift of literature radio galaxies. To gain better insights into the overall redshift evolution at $z > 2$, we combine the new faint and literature samples and bin the sources in redshift bins of $z < 2.5$, $2.5-3.0$, $3.0-3.5$, $3.5-4.0$, $4.0-5.0$, and $5.0-6.0$. The median linear sizes in each redshift bin as a function of redshift are shown in Fig. 15. The error bars on the y-axis indicate the 1σ standard deviations in the redshift bins. In the highest redshift bin, only two sources are present and therefore the small error bar is only indicative of ΔLAS .

The strong evolution of radio sizes and the size distribution of individual sources are consistent with the prediction for the evolution of radio sizes with redshift from Saxena et al. (2017), where the decrease in radio size at high redshifts is primarily dominated by IC losses. This suggests that IC scattering is the dominant energy loss mechanism at high redshifts. The enhanced IC losses, due to an increasing energy density of the cosmic microwave background radiation at high redshifts, dominate over other energy loss mechanisms such as synchrotron or adiabatic losses (see Saxena et al. 2017, for example).

5 SUMMARY

In this section, we summarize the key findings of this paper. We have obtained spectroscopic redshifts for 13 out of the 32 sources

that were selected for having an ultra-steep radio spectral index in the Saxena et al. (2018a) sample. These radio galaxies are almost an order of magnitude fainter than those that were previously discovered from large-area radio surveys and probe a new parameter space in low-radio frequency flux densities. In addition to spectroscopy, we have also obtained deep NIR imaging for a subset of the confirmed radio galaxies.

Next, we have compiled a sample of known $z > 2$ radio galaxies in the literature that have $\text{Ly}\alpha$ measurements. The literature HzRG sample was then used to compare the properties of our faint radio galaxies across redshifts. We explored the radio and optical properties of the both the faint and literature high-redshift radio galaxies. The main conclusions of this study are as follows:

(i) The newly confirmed fainter radio galaxies lie in the redshift range $0.52 \leq z \leq 5.72$, including the discovery of the highest redshift radio galaxy discovered to date, TGSS J1530+1049, which was first reported in Saxena et al. (2018b). The $\text{Ly}\alpha$ emission-line profiles show a wide variety of FWHM values, ranging from 320 to 1240 km s^{-1} . The $\text{Ly}\alpha$ luminosities and FWHM are systematically lower than what has previously been observed for brighter radio galaxies at these redshifts. Due to fainter line luminosities, narrower line widths, and no evidence of ionization by shocks in the faint HzRGs, we conclude that $\text{Ly}\alpha$ emission in these galaxies must be powered by a weak AGN.

(ii) We use K -band photometry for a subset of spectroscopically confirmed faint HzRGs presented in this paper to estimate stellar masses. Simple stellar population synthesis models are used for this. We find that the stellar masses to lie in the range $10.7 < \log M_{\star} < 11.7 M_{\odot}$, which are lower than what is seen for brighter HzRGs in the literature. Therefore, fainter HzRGs are hosted by galaxies with lower stellar masses, which are in a phase of building up their stellar mass rapidly.

(iii) We do not find any strong evidence for the presence of shocks in our sample of faint HzRGs. Therefore, we conclude that the dominant sources of ionization in faint HzRGs must either be a weak AGN or star formation. We compare the $\text{Ly}\alpha$ line luminosity and width with strong, ‘non-radio’ $\text{Ly}\alpha$ emitting galaxies and find that faint HzRGs in the redshift range $z \sim 2-4$ form a bridge between the star-forming and narrow-line AGN populations. They do not show broad-line AGN characteristics. At $z > 4$, faint HzRGs completely overlap with the star-forming galaxy population, from which we infer that they could be powered by moderate star-burst activity. It is also possible that $z > 4$ radio galaxies are powered by cold accretion, which in combination with their low stellar masses shows that they must be in the process of assembling their stellar mass.

(iv) The linear sizes of faint HzRGs do not evolve much with redshift, but the sizes of the literature HzRGs exhibit a decrease with increasing redshift. We find that the redshift evolution of sizes of sources in a combined sample of faint and literature HzRGs matches well with predictions of size evolution due to increased IC losses at high redshifts. Therefore, the dominant energy loss mechanism in radio galaxies at high redshifts is the energy loss due to interactions with the cosmic microwave background radiation, and not synchrotron or adiabatic losses.

There remain many radio sources from the original Saxena et al. (2018a) sample that still do not have a spectroscopic redshift and are high-priority targets for spectroscopy. Our new sample of HzRGs, which probes fainter flux densities over a large area on the sky, has offered glimpses into the nature of relatively less powerful radio galaxies at high redshifts. Ongoing large-area surveys that are even more sensitive at low radio frequencies, such as the LOFAR

Two-Meter Sky Survey (Shimwell et al. 2017, 2019), will uncover many more faint radio galaxies at high redshifts, furthering our understanding of how radio galaxies are fuelled and their impact on their surroundings.

ACKNOWLEDGEMENTS

We thank the referee for comments and suggestions that helped improve the quality of this work. AS wishes to thank Ricardo T. Génova Santos for fruitful discussions. AS, HJR, and KJD gratefully acknowledge support from the European Research Council under the European Unions Seventh Framework Programme (FP/2007-2013)/ERC Advanced Grant NEWCLUSTERS-321271. RAO and MM received support from CNPq (400738/2014-7, 309456/2016-9) and FAPERJ (202.876/2015). IP acknowledges funding from the INAF PRIN-SKA 2017 project 1.05.01.88.04 (FORECaST). PNB is grateful for support from STFC via grant ST/M001229/1.

This paper contains data from several different facilities. The WHT is operated on the island of La Palma by the Isaac Newton Group of Telescopes in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The HET is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maximilians-Universität München, and Georg-August-Universität Göttingen. The HET is named in honor of its principal benefactors, William P. Hobby and Robert E. Eberly. The Gemini Observatory is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina), and Ministério da Ciência, Tecnologia e Inovação (Brazil). The LBT is an international collaboration among institutions in the United States, Italy, and Germany. LBT Corporation partners are: The University of Arizona on behalf of the Arizona university system; Istituto Nazionale di Astrofisica, Italy; LBT Beteiligungsgesellschaft, Germany, representing the Max-Planck Society, the Astrophysical Institute Potsdam, and Heidelberg University; The Ohio State University, and The Research Corporation, on behalf of The University of Notre Dame, University of Minnesota, and University of Virginia.

This work has made extensive use of IPYTHON (Pérez & Granger 2007), ASTROPY (Astropy Collaboration 2013), APLPY (Robatille & Bressert 2012), MATPLOTLIB (Hunter 2007), MPDAF (Bacon et al. 2016), and TOPCAT (Taylor 2005). This work would not have been possible without the countless hours put in by members of the open-source developing community all around the world.

REFERENCES

- Afonso J. et al., 2011, *ApJ*, 743, 122
 Alam S. et al., 2015, *ApJS*, 219, 12
 Allen M. G., Dopita M. A., Tsvetanov Z. I., 1998, *ApJ*, 493, 571
 Archibald E. N., Dunlop J. S., Hughes D. H., Rawlings S., Eales S. A., Ivison R. J., 2001, *MNRAS*, 323, 417
 Arrigoni Battaia F., Hennawi J. F., Prochaska J. X., Oñorbe J., Farina E. P., Cantalupo S., Lusso E., 2019, *MNRAS*, 482, 3162
 Astropy Collaboration, 2013, *A&A*, 558, A33
 Bacon R., Piqueras L., Conseil S., Richard J., Shepherd M., 2016, MPDAF: MUSE Python Data Analysis Framework, Astrophysics Source Code Library, record ascl:1611.003
 Baum S. A., McCarthy P. J., 2000, *AJ*, 119, 2634
 Baum S. A., Heckman T. M., van Breugel W., 1992, *ApJ*, 389, 208
 Becker R. H., White R. L., Helfand D. J., 1995, *ApJ*, 450, 559
 Best P. N., Heckman T. M., 2012, *MNRAS*, 421, 1569
 Best P. N., Longair M. S., Röttgering H. J. A., 1998a, *MNRAS*, 295, 549
 Best P. N. et al., 1998b, *MNRAS*, 301, L15
 Best P. N., Röttgering H. J. A., Longair M. S., 2000, *MNRAS*, 311, 23
 Blundell K. M., Rawlings S., Eales S. A., Taylor G. B., Bradley A. D., 1998, *MNRAS*, 295, 265
 Blundell K. M., Rawlings S., Willott C. J., 1999, *AJ*, 117, 677
 Bornancini C. G., De Breuck C., de Vries W., Croft S., van Breugel W., Röttgering H., Minniti D., 2007, *MNRAS*, 378, 551
 Broderick J. W., Bryant J. J., Hunstead R. W., Sadler E. M., Murphy T., 2007, *MNRAS*, 381, 341
 Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
 Bryant J. J., Johnston H. M., Broderick J. W., Hunstead R. W., De Breuck C., Gaensler B. M., 2009, *MNRAS*, 395, 1099
 Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storchi-Bergmann T., 2000, *ApJ*, 533, 682
 Cantalupo S., Arrigoni-Battaia F., Prochaska J. X., Hennawi J. F., Madau P., 2014, *Nature*, 506, 63
 Carilli C. L., Harris D. E., Pentericci L., Röttgering H. J. A., Miley G. K., Kurk J. D., van Breugel W., 2002, *ApJ*, 567, 781
 Chabrier G., 2003, *PASP*, 115, 763
 Chambers K. C., Miley G. K., van Breugel W. J. M., 1990, *ApJ*, 363, 21
 Chambers K. C., Miley G. K., van Breugel W. J. M., Huang J.-S., 1996, *ApJS*, 106, 215
 Chambers K. C. et al., 2016, preprint (arXiv:1612.05560)
 Chonis T. S. et al., 2016, in Christopher J. E., Luc S., Hideki T., eds, Proc. SPIE Conf. Ser. Vol. 9908, Ground-based and Airborne Instrumentation for Astronomy VI. SPIE, Bellingham, p. 99084C
 Cullen F. et al., 2018, *MNRAS*, 476, 3218
 De Breuck C., van Breugel W., Röttgering H. J. A., Miley G., 2000a, *A&AS*, 143, 303
 De Breuck C., Röttgering H., Miley G., van Breugel W., Best P., 2000b, *A&A*, 362, 519
 De Breuck C. et al., 2001, *AJ*, 121, 1241
 De Breuck C., Tang Y., de Bruyn A. G., Röttgering H., van Breugel W., 2002, *A&A*, 394, 59
 De Breuck C. et al., 2010, *ApJ*, 725, 36
 Deldy P. E., Hall P. J., Schilizzi R. T., Lazio T. J. L. W., 2009, *Proc. IEEE*, 97, 1482
 Duncan K., Conselice C. J., 2015, *MNRAS*, 451, 2030
 Emonts B. H. C. et al., 2014, *MNRAS*, 438, 2898
 Feltre A., Charlot S., Gutkin J., 2016, *MNRAS*, 456, 3354
 Gabányi K. É., Frey S., Gurvits L. I., Paragi Z., Perger K., 2018, *Res. Notes Am. Astron. Soc.*, 2, 200
 Garzón F. et al., 2016, in Christopher J. E., Luc S., Hideki T., eds, Proc. SPIE Conf. Ser. Vol. 9908, Ground-based and Airborne Instrumentation for Astronomy VI. SPIE, Bellingham, p. 99081J
 Hatch N. A. et al., 2011, *MNRAS*, 410, 1537
 Hill G. J., Lilly S. J., 1991, *ApJ*, 367, 1
 Hook I. M., Jørgensen I., Allington-Smith J. R., Davies R. L., Metcalfe N., Murowinski R. G., Crampton D., 2004, *PASP*, 116, 425
 Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
 Ishwara-Chandra C. H., Sirothia S. K., Wadadekar Y., Pal S., Windhorst R., 2010, *MNRAS*, 405, 436
 Jannuzi B. T., Elston R., Schmidt G. D., Smith P. S., Stockman H. S., 1995, *ApJ*, 454, L111
 Jarvis M. J. et al., 2001a, *MNRAS*, 326, 1563
 Jarvis M. J., Rawlings S., Eales S., Blundell K. M., Bunker A. J., Croft S., McLure R. J., Willott C. J., 2001b, *MNRAS*, 326, 1585
 Jarvis M. J., Teimourian H., Simpson C., Smith D. J. B., Rawlings S., Bonfield D., 2009, *MNRAS*, 398, L83
 Kaiser C. R., Best P. N., 2007, *MNRAS*, 381, 1548
 Kaiser C. R., Dennett-Thorpe J., Alexander P., 1997, *MNRAS*, 292, 723
 Ker L. M., Best P. N., Rigby E. E., Röttgering H. J. A., Gendre M. A., 2012, *MNRAS*, 420, 2644
 Lawrence A. et al., 2007, *MNRAS*, 379, 1599
 McCarthy P. J., 1993, *ARA&A*, 31, 639
 McCarthy P. J., Baum S. A., Spinrad H., 1996, *ApJS*, 106, 281
 McLure R. J., Willott C. J., Jarvis M. J., Rawlings S., Hill G. J., Mitchell E., Dunlop J. S., Wold M., 2004, *MNRAS*, 351, 347

- McLure R. J. et al., 2018, *MNRAS*, 476, 3991
- Maddox N., 2018, *MNRAS*, 480, 5203
- Miley G., De Breuck C., 2008, *A&A*, 15, 67
- Miley G. K., Chambers K. C., van Breugel W. J. M., Macchetto F., 1992, *ApJ*, 401, L69
- Miley G. K. et al., 2004, *Nature*, 427, 47
- Miley G. K. et al., 2006, *ApJ*, 650, L29
- Morabito L. K., Harwood J. J., 2018, *MNRAS*, 480, 2726
- Nesvadba N. P. H., Lehnert M. D., De Breuck C., Gilbert A. M., van Breugel W., 2008, *A&A*, 491, 407
- Orsi Á. A., Fanidakis N., Lacey C. G., Baugh C. M., 2016, *MNRAS*, 456, 3827
- Ouchi M. et al., 2018, *PASJ*, 70, S13
- Overzier R. A. et al., 2008, *ApJ*, 673, 143
- Pentericci L. et al., 2000, *A&A*, 361, L25
- Pérez F., Granger B. E., 2007, *Comput. Sci. Eng.*, 9, 21
- Rawlings S., Eales S., Lacy M., 2001, *MNRAS*, 322, 523
- Reuland M., van Breugel W., Röttgering H., de Vries W., De Breuck C., Stern D., 2003, *ApJ*, 582, L71
- Reuland M., Röttgering H., van Breugel W., De Breuck C., 2004, *MNRAS*, 353, 377
- Rigby E. E., Best P. N., Brookes M. H., Peacock J. A., Dunlop J. S., Röttgering H. J. A., Wall J. V., Ker L., 2011, *MNRAS*, 416, 1900
- Robinson A., Binette L., Fosbury R. A. E., Tadhunter C. N., 1987, *MNRAS*, 227, 97
- Robitaille T., Bressert E., 2012, APLpy: Astronomical Plotting Library in Python, Astrophysics Source Code Library, record ascl:1208.017
- Rocca-Volmerange B., Le Borgne D., De Breuck C., Fioc M., Moy E., 2004, *A&A*, 415, 931
- Röttgering H. J. A., Lacy M., Miley G. K., Chambers K. C., Saunders R., 1994, *A&AS*, 108, 79
- Röttgering H. J. A., van Ojik R., Miley G. K., Chambers K. C., van Breugel W. J. M., de Koff S., 1997, *A&A*, 326, 505
- Röttgering H., Daddi E., Overzier R., Wilman R., 2003, *New Astron. Rev.*, 47, 309
- Saito T., Shimasaku K., Okamura S., Ouchi M., Akiyama M., Yoshida M., Ueda Y., 2008, *ApJ*, 675, 1076
- Saxena A., Röttgering H. J. A., Rigby E. E., 2017, *MNRAS*, 469, 4083
- Saxena A. et al., 2018a, *MNRAS*, 475, 5041
- Saxena A. et al., 2018b, *MNRAS*, 480, 2733
- Seifert W. et al., 2003, in Iye M., Moorwood A. F. M., eds, Proc. SPIE Conf. Ser. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes. SPIE, Bellingham, p. 962
- Seymour N. et al., 2007, *ApJS*, 171, 353
- Seymour N. et al., 2008, *ApJ*, 681, L1
- Shibuya T. et al., 2018, *PASJ*, 70, S15
- Shimwell T. W. et al., 2017, *A&A*, 598, A104
- Shimwell T. W. et al., 2019, *A&A*, 622, A1
- Singh V. et al., 2014, *A&A*, 569, A52
- Sobral D. et al., 2018, *MNRAS*, 477, 2817
- Swarup G., Ananthakrishnan S., Kapahi V. K., Rao A. P., Subrahmanya C. R., Kulkarni V. K., 1991, *Curr. Sci.*, 60, 95
- Taylor M. B., 2005, in Shopbell P., Britton M., Ebert R., eds, San Francisco, eds, ASP Conf. Ser. Vol. 347, Astronomical Data Analysis Software and Systems XIV. Astron. Soc. Pac., p. 29
- Tielens A. G. G. M., Miley G. K., Willis A. G., 1979, *A&AS*, 35, 153
- Tingay S. J. et al., 2013, *Publ. Astron. Soc. Aust.*, 30, e007
- van Breugel W., De Breuck C., Stanford S. A., Stern D., Röttgering H., Miley G., 1999, *ApJ*, 518, L61
- van Haarlem M. P. et al., 2013, *A&A*, 556, A2
- van Ojik R., Röttgering H. J. A., Carilli C. L., Miley G. K., Bremer M. N., Macchetto F., 1996, *A&A*, 313, 25
- van Ojik R., Röttgering H. J. A., Miley G. K., Hunstead R. W., 1997, *A&A*, 317, 358
- Vardoulaki E. et al., 2010, *MNRAS*, 401, 1709
- Venemans B. P. et al., 2002, *ApJ*, 569, L11
- Villar-Martín M., Tadhunter C., Clark N., 1997, *A&A*, 323, 21
- Villar-Martín M., Vernet J., di Serego Alighieri S., Fosbury R., Humphrey A., Pentericci L., 2003, *MNRAS*, 346, 273
- Villar-Martín M., Humphrey A., De Breuck C., Fosbury R., Binette L., Vernet J., 2007a, *MNRAS*, 375, 1299
- Villar-Martín M., Humphrey A., De Breuck C., Fosbury R., Binette L., Vernet J., 2007b, *MNRAS*, 375, 1299
- Willott C. J., Rawlings S., Blundell K. M., Lacy M., 1999, *MNRAS*, 309, 1017
- Willott C. J., Rawlings S., Blundell K. M., Lacy M., Hill G. J., Scott S. E., 2002, *MNRAS*, 335, 1120
- Willott C. J., Rawlings S., Jarvis M. J., Blundell K. M., 2003, *MNRAS*, 339, 173
- Wisotzki L. et al., 2016, *A&A*, 587, A98
- Wright E. L. et al., 2010, *AJ*, 140, 1868
- Wylezalek D. et al., 2013, *ApJ*, 769, 79
- Yang X.-h., Chen P.-s., Huang Y., 2015, *MNRAS*, 449, 3191
- Zirm A. W., Dey A., Dickinson M., Norman C. J., 2009, *ApJ*, 694, L31

APPENDIX A: LONG-SLIT SPECTRA

Here we show the 1D spectra (observed frame) of all confirmed radio sources observed using three different facilities. The spectra are displayed with increasing redshift, and contain information about the source name, the redshift, and the telescope used. Regions of strong skylines are marked by shaded boxes. We show a zoom-in of the Ly α emission line for radio galaxies with $z > 2$. These insets are placed on regions strongly affected by sky lines, so that the good parts of the spectrum are always visible.

The spectra have been normalized by the strength of the strongest emission line visible in the spectrum (Ly α when available). Depending on the instrument used to obtain the spectrum, the wavelength range visible varies. The ISIS instrument on WHT has two arms (Blue and Red), providing the largest wavelength range. The GMOS spectra only have the red arm. The HET spectra are split into two arms (Blue and Red), with each arm having two channels. Because LRS2 on HET is an IFU spectrograph, we could only extract the 1D spectra from those channels that had any visible continuum or emission lines. Therefore, we only show the spectrum of the channel where any emission was visible.





Figure A1. One-dimensional spectra of all spectroscopically confirmed sources with the WHT, HET, and Gemini. In the inset, we show a zoom-in of the Ly α emission line for sources with $z > 2$. The shaded boxes indicate regions of strong contamination from atmospheric sky lines. Indicated in each spectrum is the object name, redshift, and telescope code (Tel.).

Figure A1 – *continued*




Figure A1 – continued

APPENDIX B: K-BAND IMAGES

Here we show the *K*-band images for the subset of sources that were observed at NIR wavelengths. Overlaid on top of the *K*-band

image are radio contours at 1.4 GHz from Saxena et al. (2018a). The contours shown begin at 0.25 mJy, which is on average 3.5σ – 5σ ,

and are a geometric progression of $\sqrt{2}$, such that the flux density increases by a factor of 2 for every two contours.




Figure B1. K-band images taken from the LBT and GTC of the host galaxies of the radio sources, with radio contours at 1.4 GHz from Saxena et al. (2018a) overlaid on top. The contours shown begin at 0.25 mJy, which is on average 3.5σ – 5σ , and are a geometric progression of $\sqrt{2}$, such that the flux density increases by a factor of 2 for every two contours.




Figure B1 – continued

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.